The APDVS Medical Student Curriculum

The APDVS and contributors

1/31/23

Table of contents

Preface	8
Disclosures	10
Usage	11
Additional Information	12
Comments, Questions or Contributions	13
Introduction	14
Additional Vascular Surgery Educational Resources	15
Announcements & Case of the Week	17
Authors and Contributors Editors	19 20 21
Approach to Vascular Patient	22
Social Determinants of Vascular Health Introduction	24 25 25 25 26 26

Discussion Points	
Key Articles	
Additional Reading	29
Patient History and Physical Examination of the Arterial System	30
Patient History of the Arterial System	
Head and Neck History:	
Upper Extremity History:	
Abdomen Arterial History:	
Lower Extremities History:	
· ·	
Physical Exam of the Arterial System	
Head and Neck Exam:	
Upper Extremity Exam:	
Chest and Abdominal Exam:	
Lower Extremity Exam:	
Differential diagnosis of claudication	38
Carotid Disorders	41
Carotid Artery Stenosis	42
Pre/Post Questions	43
Case Based Questions	43
Operative Footage Questions	47
Introduction	49
Etiology	49
Diagnostics and Imaging	
Asymptomatic Carotid Atery Stenosis Screening	
Treatment	
Carotid Artery Endarterectomy (CEA)	
Carotid Artery Stenting (CAS)	
Optimal Medical Management	
Outcomes and Surveillance	
Asymptomatic carotid stenosis	
Symptomatic carotid stenosis	
Teaching Case	
Scenario	
Exam	
Imaging	
Discussion Points	
Key Articles	
Additional Resources	
Audible Bleeding Content	57

Websites	57
Serious Games	57
Gore Combat Manual	57
Operative Footage	58
Aortic Disorders	59
Abdominal Aortic Aneurysm (AAA)	60
Pre/Post Questions	61
Case Based Questions	61
Operative Footage Questions	64
Introduction	64
Etiology	66
Diagnostics and Imaging	66
Treatment	68
Endovascular Aortic Repair (EVAR)	69
Open Surgical Repair (OSR)	74
Outcomes	76
Surveillance	76
Teaching Case	77
Scenario	77
Exam	77
Imaging	77
Discussion Points	78
Key Articles	79
Additional Resources	80
Audible Bleeding Content	80
Websites	81
Gore Combat Manual	81
Operative Footage	81
Lower Extremity Arterial Disease	83
Claudication	84
Pre/Post Questions	85
Case Based Questions	
Introduction	
Etiology	
Anatomy	
Process of Atherosclerosis	
PAD Risk Factors	93

	Vessel Collaterization	95
	Anatomic Distribution of PAD	96
D	agnostics and Imaging	97
	Peripheral Pulse Exam	97
	Ankle Brachial Index (ABI)	97
	Pulse volume recordings (PVR)	99
	Doppler	00
	Duplex Ultrasonography	00
\mathbf{T}	reatment	
	Supervised Exercise Regimen	01
	Optimization of cardiovascular risk factors:	
	Pain Management	
Ο	utcomes	
	eaching Case	
	Scenario	
	Exam	
	Imaging	
	Discussion Points	
K	ey Articles	
	dditional Resources	
	Audible Bleeding Content	
	Gore Combat Manual	
	G ()	.05
Р	re/Post Questions	.06
	Case Based Questions	
	·	
	$troduction \dots $.11
	·	.11
\mathbf{E}	$troduction \dots $.11 .11
\mathbf{E}	troduction	11 11 12
\mathbf{E}	troduction	11 11 12
\mathbf{E}	troduction	.11 .12 .13 .13
\mathbf{E}	troduction	.11 .12 .13 .13
\mathbf{E}	troduction	.11 .12 .13 .13
E D	troduction	.11 .12 .13 .13 .13
E D	troduction 1 ciology 1 tagnostics and Imaging 1 Patient History 1 Physical Exam 1 Non-invasive Hemodynamic Testing 1 Angiography 1 Classification Systems 1	.11 .12 .13 .13 .14 .15
E D	troduction	.11 .12 .13 .13 .13 .14 .15
E D	troduction	.11 .12 .13 .13 .13 .14 .15
E D	troduction	.11 .12 .13 .13 .14 .15 .19
E D	troduction	.11 .12 .13 .13 .14 .15 .19 .19 .21
E D	troduction	.11 .12 .13 .13 .14 .15 .19 .19 .21

Teaching Case				 	 	 				123
Scenario				 	 	 				123
Exam .				 	 	 				124
Imaging				 	 	 				124
	n Points									
Key Articles .				 	 	 				125
Additional Res	ources			 	 	 				125
Audible E	Bleeding Content			 	 	 				125
Gore Con	nbat Manual			 	 	 				126
Operative	e Footage			 	 	 	 •	 •	 •	126
Miscellaneous T	opics									127
Basics of Endovas	•		•							128
	tions									
	ed Questions									
	omy and Vascular									
	d Guided Access									
,	ers, and Sheaths									
,	ts, and Endografts									
0	ts									
	vascular Procedur									
	r Placement									
	tremity Angiogran									
	ams									
Access Co	omplications			 	 	 				151
Endovasc	ular Surgical Com	plications		 	 	 				154
Teaching Case				 	 	 				155
Scenario				 	 	 				155
Exam .				 	 	 				156
Discussion	n Points			 	 	 				156
Key Articles .				 	 	 				157
Additional Res	ources			 	 	 				157
Audible E	Bleeding Content			 	 	 				157
Websites										158

	Fore Combat Manual	
Anonyr	us Feedback	160
Referer	es s	161

Preface



Association of Program Directors in Vascular Surgery

This content was developed as part of the Association of Program Directors in Vascular Surgery's (APDVS) medical student curriculum. Each chapter covers a key domain of vascular surgery pathology and treatment and is associated with an Audible Bleeding episode which you can access from the link embedded in the text.

This eBook would not have been possible without the Association of Program Directors in Vascular Surgery (APDVS), the guidance and resources of Dr. Sharif Ellozy and Dr. Adam Johnson, the Audible Bleeding Vascular Surgery Exam Prep project, and the Audible Bleeding Team. This project is the direct result of Dr. Chelsea Dorsey et al.'s medical education research (see below) and the work of the Society of Vascular Surgery's Resident Student Outreach Committee (SVS RSOC).

Below, please see the two papers to understand the medical education need addressed by this eBook.

- Vascular Surgery Curriculum for Medical Students: A National Targeted Needs Assessment
- The Value of a Vascular Surgery Curriculum for Clinical Medical Students: Results of a National Survey of Nonvascular Educators

Editors: The Association of Program Directors in Vascular Surgery and Ezra Schwartz.

Version 01.02

Disclosures

This is an publication of the APDVS and therefore we share the same conflicts of interest. That noted, this eBook is the product of voluntarily donated time. This work has no financial backing and is not-for-profit. We are not receiving any funding or benefits from outside resources referenced. We include resources we believe will benefit you, the learner.

Usage

This eBook is intended to be a medical student level, easily accessible review for vascular surgery rotations and general medical education. This eBook is paired with slide decks, preand post-reading questions and a teaching case produced by the APDVS in addition to podcast content produced by Audible Bleeding, a publication of the Society of Vascular Surgery.

We are also excited to include an open source annotations software called hypothes.is. By creating a Hypothes.is account, you will be able to creates notes in the eBook as you read. You can choose to make your notes private or public. As this eBook is a community-led initiative, please consider making your notes public if you feel they would benefit your peers or the editors of the book. Given the public nature of annotations, if you choose to post public notes, these annotations are regarded as contributions to the eBook and we expect readers and listeners to follow a Contributor Code of Conduct. We expect all participants, ourselves included, to maintain a safe space and behave professionally.

Thanks to the flexibility and accessibility afforded as an eBook, we can (and will) update the book with extreme ease. Updates may include but are not limited to: slide decks, new preand post-reading questions, and relevant Audible Bleeding podcast episodes. Please see the announcements page to stay up to date on developments.

This eBook is free to the public and licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 United States License.

Additional Information

Please consider becoming a member of the Society of Vascular Surgery. Benefits include free access to the Journal of Vascular Surgery, mentorship and scholarship opportunities, and more! A variety of membership types exist, including those for medical students, general surgery residents, and integrated vascular trainees.

Visit the ChooseVascular website to learn why we chose vascular surgery and determine if you wish to choose vascular too! This site is complete with a student resources page, and information to pair you with a vascular surgery mentor or a vascular surgery interest group, this website is a wonderful introduction the work and community of vascular surgery.

Comments, Questions or Contributions

Please visit our github page or send the APDVS an email.

This book is built on Quarto.

Introduction

Additional Vascular Surgery Educational Resources

We hope you use this eBook as a review for your rotations, undergraduate medical licencing examinations, and entry to independent practice. This resource is a summary and is by no means comprehensive. That noted, in every chapter we have included links to additional resources if you wish to gain a more comprehensive understanding of the topics covered. As our mission is to make vascular surgery education accessible, we prioritize open access publications and free materials.

A few highly recommended additional resources include:

- The Gore Medical Vascular and Endovascular Surgery Combat Manual is an informative and entertaining read intended as a vascular surgery crash course for medical students, residents, and fellows alike. Highly accessible with a thoughtfully determined level of detail, but lacking in learning activities (e.g. questions, videos, etc.), this resource is a wonderful complement to the APDVS eBook.
- TeachMe Surgery is a student friendly online resource complete with short quizzes to help cement learning and recommended readings from the literature. Clear and concise, this is a great just-in-time learning resource.
- The Audible Bleeding Medical Student Archive contains podcast episodes tailored for medical students. We suggest paying special attention to the *Holding Pressure* series. Whereas this eBook is organized by disease, *Holding Pressure* episodes are organized by surgical procedure. These episodes contain a wealth of high-yield information and we will include links to *Holding Pressure* episodes throughout the eBook when applicable.
 - You can listen to Audible Bleeding episodes on Apple Podcasts, Spotify, Google Podcasts, Stitcher, or Soundcloud
- The Audible Bleeding Exam Prep curriculum and the associated podcast series. While this resource is intended for postgraduate trainees and fellows and therefore beyond the scope of this curriculum, it is a well authored, no-nonsense resource for those who wish to deepen their knowledge base. The accompanying podcast series also allows you to learn or study on the go!

- You can listen to Audible Bleeding episodes on Apple Podcasts, Spotify, Google Podcasts, Stitcher, or Soundcloud
- Vascular Training.org is an online repository of vascular surgery trainee resources compiled during the stages of the COVID-19 pandemic when providing traditional education was challenging and learning content was increasingly available online. Find videos, landmark and other informative papers, and device information here. Like the Audible Bleeding Exam Prep curriculum, the primary audience for this content is postgraduate trainees but is very accessible for undergraduate students who wish to deepen their knowledge base. This resource is password protected. The password is: Vasceducation 2021!

Announcements & Case of the Week



Please return to this page for updates on website, podcast, and activity developments.

eBook Chapters

Claudication chapter hot off the press!

We are starting to post our very own illustrations!

Chapters in development include: - Patient History and Physical Examination of the Venous System - Peripheral artery disease (PAD) including topics such as acute limb ischemia (ALI) and chronic limb threatening ischemia (CLTI). - Aortic dissection - Thoracic outlet syndrome - Endovascular surgery basics - Vascular lab primer

Podcast Episodes

The next podcast in development is an introduction to the APDVS curriculum.

Case of the Week

To harness the learning principles of spaced practice, retrieval practice, interleaving and concrete examples, we invite students and residents to write a case of the week and submit it to APDVS for review.

Please try to build the case using material covered in the eBook. We ask you to write for the medical student level. Please include discussions to questions including why the correct is answer is correct and why the incorrect answers are incorrect.

Once reviewed, we will post it here for the community! We hope you will take part in this mini-publication exercise!

Please submit your case to APDVSebook@vascularsociety.org.

Authors and Contributors

This eBook is the result of a partnership between the APDVS and the Audible Bleeding Podcast, each of which is comprised of hard working individuals dedicated to medical education and the promotion of vascular surgery. The authors and contributors of this eBook have freely donated their time and experience to create this resource. Please follow them on twitter or other social media platforms to return this debt of gratitude and to, as Audible Bleeding says, "keep a finger on the pulse."

Editors

Ezra Schwartz, MDCM, MS, MMS. Harvard Medical School @EzraSchwartz10
Yasong Yu, MD. Rutgers New Jersey Medical School @YasongYuMD
Chaitanya Dandu, MD. Yale School of Medicine @ChaituDandu

Faculty Authors

Jonathan Bath, MD, FACS, DFSVS. Associate Professor of Surgery; Program Director, Vascular Surgery Fellowship; *University of Missouri School of Medicine* Academic Profile

Cassius Iyad Ochoa Chaar, MD, MS, RPVI. Associate Professor of Surgery; Yale School of Medicine Academic Profile @MdChaar

Dawn Coleman, MD, FACS. Chief, Division of Vascular and Endovascular Surgery; *Duke University School of Medicine* Academic Profile @ColemanDM_vasc

Sharif Ellozy, MD. G. Tom Shires Faculty Scholar in Surgery, Associate Professor of Clinical Surgery, Program Director of the Vascular Surgery Fellowship; Weill Cornell Medical College Academic Profile@SharifEllozy

Erin Greenleaf, MD, MS, RPVI Assistant Professor; Baylor College of Medicine Academic Profile

Angela Kokkosis, MD, FACS. Associate Professor of Surgery; Program Director, Vascular and Endovascular Residency; Director, Carotid Interventions; Co-Medical Director, Non-Invasive Vascular Laboratory; Stony Brook University Academic Profile @Angela_Kokkosis

Loay S Kabbani, MD, MHSA, FACS. Program Director, Vascular Surgery Fellowship; Henry Ford Health Academic Profile @loaykabbani

Michael Malinowski, MD, MEHP, FACS. Associate Professor; Medical College of Wisconsin, Froedert Hospital Academic Profile

Trainee Authors

Marvin Chau, MD. Integrated Vascular Surgery Resident; UCLA Health Academic Profile@MChau_MD

Nakia Sarad, DO, MS. General Surgery Resident, Vascular Surgery Research Fellow; Weill Cornell Medicine Academic Profile @NakSaradDO

Sree Kannathasan, MD. Integrated Vascular Surgery Resident; McGill University @VulnerableAorta

Regis Hila, MD. General Surgery Resident; Northwell Health @HilaRegis

Emily Goodman, MD. Vascular Surgery Fellow; The University of Chicago Academic Profile @emily_goodmanMD

Raj S. Lingnurkar, MD. General Surgery Resident; Trinity Health Mid-Atlantic @rslingnurkar

Joseph Yoo, MD. Integrated Vascular Surgery Resident; Jefferson Health / Einstein Health-care Network Academic Profile

Abena Appah-Sampong, MD. General Surgery Resident; Brigham and Women's Hospital @AbenaSamp

Student Contributors

 $\textbf{Ivana Zhang} \ \text{MS4}; \ \textit{Jacobs School of Medicine And Biomedical Sciences at University at Buffalo}$

 ${\bf Shannon\ McDonnell\ MS4};\ Stritch\ School\ of\ Medicine\ at\ Loyola\ University\ Chicago$

Medical Illustrators

Ivana Zhang MS4; Jacobs School Of Medicine And Biomedical Sciences at University at Buffalo

Approach to Vascular Patient

Social Determinants of Vascular Health

Abena Appah-Sampong, MD, Ezra Schwartz, MD, CM, MS, Sharif Ellozy, MD.

Note

By the end of this chapter, students will:

- Review terms important to the discussion about social determinants of health.
- Be able to predict relevant questions to ask patients when obtaining a social history.
- Relate the importance of a thorough social history to the incidence, morbidity, and mortality of vascular disease.
- Recognize opportunities for one's biases to impact quality of care.

Key Facts

- Racial and ethnic minorities encounter social determinants of health and systemic racism that exacerbate the incidence, prevalence, and outcomes of many vascular diseases.
- 2. Addressing the challenges of healthcare deserts, particularly in rural areas, is vital for ensuring equitable access to vascular surgery care.
- 3. The underrepresentation of women in vascular surgery clinical trials highlights the imperative for concerted efforts to address gender disparities, enhance inclusion, and ensure findings are generalizable to broader populations.
- 4. Many veterans face socioeconomic, systemic, and mental health challenges stemming from their military service that challenge the management of chronic and complex vascular diseases.

Introduction

"Do we not always find the diseases of the populace traceable to defects in society?" - Rudolf Virchow

Multiple studies have demonstrated significant disparities exist across the spectrum of vascular disease including prevalence, treatment, and disease outcomes. Furthermore, it has been demonstrated that social determinants of health (SDoH), comprising multiple factors including patient socioeconomic status, race or ethnicity, gender, and geographic location, can have potent effects on health outcomes. There are some estimates that SDoH account for 80-90 percent of the modifiable contributors to healthy outcomes for a population, and must be addressed in order to rectify persistent healthcare disparities. (Remington, Catlin, and Gennuso 2015)

Vascular surgeons treat patients with advanced atherosclerosis and diabetes, which are diseases that disproportionately affect vulnerable and socioeconomically disadvantaged patients. Given this reality, it is important for vascular surgeons to have a strong foundational understanding of the social determinants of health and health disparities that lead to unequal care among patients. This textbook chapter does not intend to be comprehensive, but rather an introduction to scope of health disparities in vascular surgery.

Definitions

Health Disparities: The preventable differences in the burden of disease, injury, violence, or opportunities to achieve optimal health that are experienced by socially disadvantaged populations. ("Health Disparities | DASH | CDC" 2023)

Social Determinants of Health: The circumstances in which people are born, grow up, live, work, age, and the systems put in place to deal with illness. ("Social Determinants of Health: Key Concepts," n.d.)

Structural Racism: A system in which public policies, institutional practices, cultural representations, and other norms work in various, often reinforcing ways to perpetuate racial group inequity. (Allison et al. 2023)

Health Disparities in Vascular Surgery

Race, Ethnicity, and Structural Racism

There is an abundance of literature documenting racial and ethnic disparities across the spectrum of vascular disease including thoracic and aortic aneurysms, carotid disease, and hemodialysis access. However, disparities spanning the disease continuum of peripheral artery disease (PAD) is one of the most robustly studied areas in surgery disparities research.

A scientific statement published by the American Heart Association outlines the scope of health disparities in PAD. Looking at prevalence, it has been demonstrated Black patients have a higher prevalence of PAD, even after controlling for traditional cardiovascular risk factors. Prevalence of PAD among other racial and ethnic groups is less documented, but limited evidence suggests higher rates of PAD among Cuban populations especially when compared to other Hispanic and Latinx groups. Risk factor control and preventative measures are important management tools to prevent the progression of PAD to critical limb threatening ischemia (CLTI) and limb loss. However, Black and Hispanic patients are more likely to present with hypercholesterolemia but less likely to be prescribed an aspirin or statin. (Allison et al. 2023)

Looking at PAD outcomes, studies have found that patients identifying as Black have nearly twofold greater rate of leg amputation, even after controlling for relevant confounders. Additionally, Black patients who undergo amputation are less likely to have any limb-related admission, suggesting less aggressive limb salvage among this population. Higher rates of diabetes and deep soft tissue infection are strongly correlated with the higher burden of amputation in Black patients. (Hackler, Hamburg, and White Solaru 2021)

Due to structural racism, racial and ethnic minority patients are often disproportionately burdened with socioeconomic barriers. One study evaluating the etiology of health care access among Hispanic patients found that Hispanic patients were more likely to delay medical care for socioeconomic reasons including inability to afford costs, time away from employment, and childcare issues. (Bazikian et al. 2023)

Access and Care Delivery:

Millions of Americans live in communities considered 'healthcare deserts,' where they lack adequate access to important healthcare services including primary care, hospitals, emergency services, pharmacies, trauma centers, and low-cost health centers. ("Mapping Healthcare Deserts: 80," n.d.) Many healthcare deserts are often concentrated in rural locations highlighting the unique geographical challenges that these patients face. Furthermore, certain vascular diseases including chronic limb threatening ischemia (CLTI) require specialized, multidisciplinary care for optimal management, but many communities are considered "vascular deserts" in which there is inadequate access to expert vascular surgery care. (DiLosa et al. 2023) Addressing the

challenges posed by healthcare and vascular deserts, particularly in rural areas, is crucial for ensuring equitable access to essential services. Continued investigation and investment into these issues are imperative to develop effective solutions that can bridge the healthcare gaps and provide optimal care.

Sexual and Gender Minorities:

A growing body of research explores the impact of sex and gender on various aspects of vascular disease. A systematic review and meta-analysis of 236 studies demonstrates an association between female sex or gender and poorer outcomes in individuals with Abdominal Aortic Aneurysm (AAA) and Peripheral Artery Disease (PAD). (Lee et al. 2022) It has been demonstrated that women have higher risk of AAA rupture, prolonged lengths of stay, and decreased survival following both elective and emergent AAA repair. (Dillavou, Muluk, and Makaroun 2006) The lack of representation of women in large-scale vascular surgery clinical trials has also become an area of increasing concern, given the pivotal role these studies play in shaping the standard of care and management guidelines. Despite efforts to enhance women's participation in clinical trials, evidence suggests persistent underrepresentation in randomized control trials, potentially limiting the broader applicability of their findings. (Hoel et al. 2009) The multifactorial nature of this underrepresentation, influenced by factors like patient preference, bias, and local demographics, necessitates sustained efforts to achieve diverse recruitment and ensure a more comprehensive understanding of vascular disease across diverse patient populations.

Veteran Populations:

Military veterans represent a unique population with distinct healthcare needs. While veterans may benefit from the ability to receive care through the Department of Veterans Affairs (VA) healthcare system, there remain notable challenges and disparities for this population. Barriers to medical treatment in the veteran population broadly include long wait times at VA facilities, mental health illness and stress related to previous deployments, challenges reintegrating into civilian life, and unmet basic needs such as housing and employment. (Misra-Hebert et al. 2015) Because of these barriers, veteran populations often suffer higher incidence of medical comorbidities compared to civilian population. Assari (2014) Looking at vascular pathologies, a large 2017 epidemiological study demonstrated that there is a higher incidence of PAD and associated mortality among veterans, with an estimated 30% of veterans dying within 3.8 years of a PAD diagnosis. (Willey et al. 2018) Additional studies have shown that demonstrated that mental illness may compound vascular comorbidities within the veteran population, as veterans suffering from comorbid depression face significantly higher risk of amputation and death from PAD. (Arya et al. 2018) Ultimately, understanding the complex social determinants and healthcare needs of veterans is essential for developing targeted interventions and support systems to improve the health and well-being of this vulnerable population.

Teaching Case

Scenario

A 65 year old man is brought to the emergency room with altered mental status, fevers, and severe pain, swelling, and erythema in his right foot. On exam, the foot is warm to the touch and you observe a deep, ulcerating wound with surrounding necrotic tissue. His labs are notable for an elevated white count and HbA1c of 10. His daughter, who is the patient's legal medical decision maker, is at the patient's bedside and tells you that the patient was laid off from work 6-months ago and since then has not been able to afford insulin to manage his diabetes. You call your attending surgeon about the patient's concerning findings and recommend an emergency operation. Your attending sighs and says "I'll never understand what kind of person lets their disease get this bad."

Case inspired by University of Michigan Cultural Complications Curriculum

Discussion Points

N.B. There is no prepared answers for the questions below.

However, we feel this chapter contains all the necessary information to answer the questions. If not, please let us know!

- 1. How do socioeconomic factors impact a patient's ability to manage chronic conditions?
- 2. In this scenario, how might the attending surgeon's assumptions affect the quality of care provided?
- 3. Do surgeons have a responsibility in addressing underlying socioeconomic challenges that affect a patient's health?

Key Articles

- 1. Remington PL, Catlin BB, Gennuso KP. The County Health Rankings: rationale and methods. Popul Health Metr. 2015;13(1):11. doi:10.1186/s12963-015-0044-2.(Remington, Catlin, and Gennuso 2015)
- 2. Health Disparities | DASH | CDC. Published May 26, 2023. Accessed December 7, 2023. https://www.cdc.gov/healthyyouth/disparities/index.htm ("Health Disparities | DASH | CDC" 2023)
- 3. Social determinants of health: Key concepts. Accessed December 4, 2023. https://www.who.int/news-room/questions-and-answers/item/social-determinants-of-health-key-concepts ("Social Determinants of Health: Key Concepts," n.d.)
- 4. Allison MA, Armstrong DG, Goodney PP, et al. Health Disparities in Peripheral Artery Disease: A Scientific Statement From the American Heart Association. Circulation. 2023;148(3):286-296. doi:10.1161/CIR.000000000001153.(Allison et al. 2023)
- Hackler EL, Hamburg NM, White Solaru KT. Racial and Ethnic Disparities in Peripheral Artery Disease. Circ Res. 2021;128(12):1913-1926. doi:10.1161/CIRCRESAHA.121.318243.(Hackler, Hamburg, and White Solaru 2021)
- 6. Mapping Healthcare Deserts: 80% of the Country Lacks Adequate Access to Healthcare GoodRx. Accessed December 8, 2023. https://www.goodrx.com/healthcare-access/research/healthcare-deserts-80-percent-of-country-lacks-adequate-healthcare-access.("Mapping Healthcare Deserts: 80," n.d.)
- 7. DiLosa KL, Nguyen RK, Brown C, Waugh A, Humphries MD. Defining Vascular Deserts to Describe Access to Care and Identify Sites for Targeted Limb Preservation Outreach. Ann Vasc Surg. 2023;95:125-132. doi:10.1016/j.avsg.2023.05.025.(DiLosa et al. 2023)
- 8. Lee MH, Li PY, Li B, Shakespeare A, Samarasinghe Y, Feridooni T, Cuen-Ojeda C, Alshabanah L, Kishibe T, Al-Omran M. A systematic review and meta-analysis of sex- and gender-based differences in presentation severity and outcomes in adults undergoing major vascular surgery. J Vasc Surg. 2022 Aug;76(2):581-594.e25. doi: 10.1016/j.jvs.2022.02.030. Epub 2022 Mar 5. PMID: 35257798.(Lee et al. 2022)
- 9. Dillavou ED, Muluk SC, Makaroun MS. A decade of change in abdominal aortic aneurysm repair in the United States: Have we improved outcomes equally between men and women? J Vasc Surg. 2006;43(2):230-238. doi:10.1016/j.jvs.2005.09.043.(Dillavou, Muluk, and Makaroun 2006)
- 10. Hoel AW, Kayssi A, Brahmanandam S, Belkin M, Conte MS, Nguyen LL. Under-representation of women and ethnic minorities in vascular surgery randomized controlled trials. J Vasc Surg. 2009;50(2):349-354. doi:10.1016/j.jvs.2009.01.012.(Hoel et al. 2009)

- Misra-Hebert AD, Santurri L, DeChant R, Watts B, Rothberg M, Sehgal AR, Aron DC. Understanding the Health Needs and Barriers to Seeking Health Care of Veteran Students in the Community. South Med J. 2015 Aug;108(8):488-93. doi: 10.14423/SMJ.00000000000000326. PMID: 26280777; PMCID: PMC4544768. (Misra-Hebert et al. 2015)
- 12. Kazis LE, Miller DR, Clark J, Skinner K, Lee A, Rogers W, Spiro A 3rd, Payne S, Fincke G, Selim A, Linzer M. Health-related quality of life in patients served by the Department of Veterans Affairs: results from the Veterans Health Study. Arch Intern Med. 1998 Mar 23;158(6):626-32. doi: 10.1001/archinte.158.6.626. PMID: 9521227.(Kazis et al. 1998)
- 13. Assari S. Veterans and risk of heart disease in the United States: a cohort with 20 years of follow up. Int J Prev Med. 2014 Jun;5(6):703-9. PMID: 25013689; PMCID: PMC4085922. (Assari 2014)
- 14. Willey J, Mentias A, Vaughan-Sarrazin M, McCoy K, Rosenthal G, Girotra S. Epidemiology of lower extremity peripheral artery disease in veterans. J Vasc Surg. 2018 Aug;68(2):527-535.e5. doi: 10.1016/j.jvs.2017.11.083. Epub 2018 Mar 24. PMID: 29588132; PMCID: PMC6132057.(Willey et al. 2018)
- 15. Arya S, Lee S, Zahner GJ, Cohen BE, Hiramoto J, Wolkowitz OM, Khakharia A, Binney ZO, Grenon SM. The association of comorbid depression with mortality and amputation in veterans with peripheral artery disease. J Vasc Surg. 2018 Aug;68(2):536-545.e2. doi: 10.1016/j.jvs.2017.10.092. Epub 2018 Mar 24. PMID: 29588133; PMCID: PMC6057818.(Arya et al. 2018)
- 16. Bazikian S, Urbina D, Hsu CH, Gonzalez KA, Rosario ER, Chu DI, Tsui J, Tan TW. Examining health care access disparities in Hispanic populations with peripheral artery disease and diabetes. Vasc Med. 2023 Dec;28(6):547-553. doi: 10.1177/1358863X231191546. Epub 2023 Aug 29. Erratum in: Vasc Med. 2024 Feb;29(1):NP1. PMID: 37642640; PMCID: PMC10712238. (Bazikian et al. 2023)

Additional Reading

- University of Washington Chapter on Taking a Social History
- University of Michigan Cultural Complications Curriculum

Patient History and Physical Examination of the Arterial System

Loay S Kabbani, MD, MHSA, FACS and Cassius Iyad Ochoa Chaar, MD, MS, RPVI

Listen to your patient, [they are] telling you the diagnosis. - William Osler.

Specialized testing should always be guided by the initial clinical impressions.

Though the electronic medical record helps eliminate repetitive questions and tests, it tends to divert attention from the importance of talking to and examining the patient. Interacting with the patient and family during the history and physical is extremely important to **build** rapport with the patients and their family.

For vascular patients passing the "eyeball test" is important before discussing complex procedures. This includes knowing the level of independent living.

Patient History of the Arterial System

For vascular specialists, a focused arterial history should include the arteries of the neck, torso, and extremities. Many diseases are systemic. For example, atherosclerotic disease affects the carotid arteries, heart, upper extremities, abdominal blood vessels, and lower extremities. Patients with an abdominal aortic aneurysm may have synchronous peripheral aneurysms.

Head and Neck History:

- The history should focus on the carotid artery disease—see ?@sec-carotidarterystenosis for more detail—specifically, any history of stroke, transient ischemic attacks (TIA), or amaurosis fugax.
- Amaurosis fugax is described as a curtain or shade causing transient monocular blindness. This is usually caused by emboli from carotid disease on the ipsilateral side but may be caused by other embolic sources, migraine, or giant cell arteritis.
- Syncopal episodes are usually not vascular in nature. However, in patients with subclavian stenosis, syncope may be a manifestation of subclavian steal syndrome.
- History of radiation to neck, prior carotid interventions or prior surgery to the neck should be elucidated and prior studies should be obtained.

Upper Extremity History:

- Evaluation of the upper extremities should focus on the timing (acute vs. chronic) and degree of ischemia (claudication vs. chronic limb-threatening ischemia).
- Acute limb ischemia (ALI) is a sudden occlusion of the blood supply with no time for collateral vessels to develop. It is characterized by the 6 Ps. Pulselessness, pain, pallor, paresthesia, poikilothermia (cold), and paralysis. The most common etiology of acute limb ischemia is embolic. However, uncommon causes of ALI in the upper extremity may include thrombosis of a subclavian–axillary aneurysm.
- Intermittent hand coldness associated with pain and numbness may reflect Raynaud's syndrome (associated with cold exposure) or vasospam of small vessels due to conditions like frostbite or scleroderma.
- Raynaud's phenomenon presents as episodes of vasospasm in response to cold or stress. During an attack, the patient describes the affected areas turning white, blue, and red associated with feeling cold and numb. As the circulation improves and the affected areas turn red, the patient may experience throbbing, tingling, and swelling.
- Episodes of ischemia to the digits may be persistent, severe and associated with underlying obliterative microangiopathy and manifest with pain and tissue loss (ulceration or gangrene).
- Chronic ischemia of the upper extremity is uncommon and can manifest with arm claudication (exertional fatigue). The etiology is most commonly proximal obstruction due to atherosclerotic disease in the subclavian artery. Uncommon causes can include Takayasu arteritis and thoracic outlet syndrome (TOS).
- Exertion of the arm causing posterior cerebral circulation symptoms (diplopia, dysarthria, dizziness, drop attacks, vertigo, syncope, and ataxia) may reflect a subclavian steal syndrome caused by subclavian artery stenosis proximal to the vertebral artery.

Abdomen Arterial History:

- A majority of vascular pathology in the abdomen can be attributed to aneurysmal disease
 or atherosclerotic disease. Other etiologies of abdominal vascular diseases can include
 compression syndromes, embolisms, or dissections.
- Aneurysmal disease in the abdomen is usually asymptomatic.
- Patients with known abdominal aortic aneurysm (AAA) who present with flank or back pain should be presumed to have a ruptured aneurysm until proven otherwise.
- On the other hand, mesenteric atherosclerosis can present with a constellation of symptoms. The typical trifecta of symptoms is weight loss, sitophobia (fear of food), and postprandial pain. Atherosclerosis of the renal arteries, known as renal artery stenosis, may present as severe hypertension, especially in young adults.
- Aortoiliac occlusive disease may present with Leriche's syndrome. **Leriche's syndrome** is defined as bilateral hip and buttock claudication, absent femoral pulses, and impotence.
- Vascular etiologies of acute abdominal pain include ruptured AAA aneurysm, which
 presents as severe pain radiating to the back or acute mesenteric ischemia, which presents
 as pain out of proportion to physical exam

Lower Extremities History:

- An evaluation of the lower extremities for peripheral artery disease should focus on the timing (acute vs. chronic) and degree of ischemia (claudication vs. chronic limbthreatening ischemia).
- Acute limb ischemia (ALI) is the sudden occlusion of the blood supply to a peripheral limb. It is typically characterized by the 6 Ps. Pulselessness, pain, pallor, paresthesia, poikilothermia (cold), and paralysis.
- Claudication is defined as cramping pain in the leg induced by exercise and relieved by rest. It occurs after a fixed and reproducible distance and resolves with rest. The pain is described as discomfort, cramping, numbness, or tiredness in the legs. Claudication most commonly occurs in the calf muscles, but it can also affect the feet, thighs, hips, and buttocks. It is crucial to determine to what extent the patient's claudication affects their lifestyle during history taking.
- Table 1 lists various differential diagnosis of leg pain, which can help differentiate claudication from pseudoclaudication and other types of leg pain. Table 1 is also helpful in differentiating between other types of leg pain, pseudoclaudication, and true claudication.
- Chronic limb-threatening ischemia (CLTI) is when a patient with PAD has rest pain or tissue loss. CLTI manifests as rest pain or tissue loss. Rest pain is characterized as pain in the dorsum of the foot and toes at rest (i.e., without exertion such as walking). The pain may worsen with leg elevation and many patients wake up in the middle of the

- due the pain. They often state the pain improves by dangling their affected limb over the side of their bed, which leads to better blood flow to the foot thanks to gravity.
- Tissue loss is often present in the form of a nonhealing wound, ulcer, or gangrene anywhere along the toes, foot, or lower leg. It is imperative you determine how long the tissue loss has been present and if it has been improving. Often, patients with CLTI have multiple levels of arterial disease and are at a much higher risk of amputation compared to patients with simple claudication.

Physical Exam of the Arterial System

The arterial system is extensive, and a proper vascular exam should include the entire arterial system.

The exam starts with the nurse checking the vital signs and bilateral upper extremity blood pressures. A difference of > 10 mmHg between the upper extremities may indicate significant hemodynamic stenosis.

i Key Notes

Arterial Bruit: Normal arterial flow is laminar and therefore silent. However, when blood flow becomes turbulent it can often be heard (bruit) or palpated (thrill). Although murmur and bruit are synonymous, audible turbulence originating from the heart is called a murmur. Outside of the heart, audible turbulence is referred to as a bruit. The presence of a bruit does not necessarily indicate significant arterial stenosis.

Pulses are described as:

- 0 or absent
- 1+ or diminished
- 2+ or normal
- 3+ or prominent or aneurysmal.
- 4+ or bounding or aneurysmal.

In patients with weak pulses, the examiner must take care not to confuse the patient's pulse with involuntary muscle twitches or their own pulse.

When a pulse is not palpated, a Doppler probe is used to assess the blood flow. The doppler signal can be triphasic, or biphasic, or monophasic.

Doppler Signals:

• Triphasic and biphasic doppler signals indicate good blood flow.

- Monophasic signal correlates with moderate to severe decrease in arterial blood flow.
- In severe ischemia, a soft continuous venous signal may be all that is heard.

Listen to examples of Doppler signals here.

On Exam:

- Temperature changes may help demarcate the level of disease.
- Sensory loss may be present in acute ischemia, chronic limb threatening ischemia, and chronic neuropathy.

Head and Neck Exam:

Inspection:

- Pulsatile masses in the neck are usually tortuous carotid arteries mistaken for carotid aneurysms. Carotid aneurysms are usually near the carotid bifurcation, while tortuous carotid arteries are usually at the base of the neck. Carotid body tumors are also at the carotid bifurcation. Both carotid body tumors and aneurysms are not visible until they are large.
- If the patient complains of amaurosis fugax (i.e., transient vision loss), fundoscopy may reveal cholesterol plaques called Hollenhorst plaques. These are thought to originate from the carotid plaque.

Palpation:

- The carotid pulse is palpated on the medial border of the sternocleidomastoid muscle. Carotid palpation is generally not performed routinely as it may cause a syncopal episode in elderly patients with sensitive carotid bulbs.
- A robust temporal pulse anterior to the ear is a sign of a patent common and external carotid artery sign.
- A large supraclavicular pulse may indicate an enlarged subclavian artery. Otherwise, the subclavian artery is usually not palpable.

Auscultation:

• Auscultate the carotid arteries for bruits.

- Using the stethoscope bell, you can hear the S1, and S2 heart sounds in the carotid artery in the mid-neck. A bruit heard in the neck is not normal. This could be transmitted from the heart or could be from a kink or narrowing in the carotid artery. The carotid bruit is loudest in the mid neck over the carotid bifurcation. A heart murmur is loudest in the upper chest.
- The intensity of the bruit and pitch do not correlate with the severity of stenosis. A tight stenosis may have low flow and thus a faint bruit.
- When a carotid bruit is heard, only 25% will have significant stenosis (75% or greater), and 50% will not have any stenosis.

Upper Extremity Exam:

Inspection

- Pink fingertips with capillary refill times < 2 seconds are a reliable sign of adequate perfusion. Ischemia in the extremities manifests as paleness with poor to no capillary refill. Chronic ischemia manifests with muscle atrophy.
- Raynaud phenomenon is characterized by a sharply demarcated triphasic color change after exposure to cold or emotional stress. First, the capillaries contract after the stressor, causing a characteristic white appearance. Then, as the capillaries open a little, deoxygenated blood re-perfuses sluggishly which leads to a hypoxic blue color of the distal extremity. Finally, when the capillaries recover and hyper-dilate, the affected limb becomes red and hyperemic. Raynaud's phenomena may occur idiopathically (Raynaud's Disease) or secondarily due to autoimmune disease. It is often provoked by emotional distress or exposure to the cold.

Palpation

- Palpate the axillary artery in the upper arm in the groove between the biceps and triceps muscle.
- Palpate the brachial artery in the antecubital fossa just medial to the biceps tendon.
- Palpate the radial artery on the wrist's flexor surface just medial to the radial styloid.
- Palpate the ulnar artery on the wrist's flexor surface just medial to the distal ulna; it lies deeper than the radial artery and may not be palpable.
- Absent pulses should initiate a search for a cause such as proximal atherosclerotic stenosis
 in older adults or autoimmune disease such as Takayasu's in young females.
- Aneurysm of the subclavian artery and axillary artery (assessed above and below the clavicle) are difficult to palpate if small. Brachial artery aneurysms are usually pseudoaneurysms from trauma or arterial access. Ulnar artery aneurysm occurs from repetitive trauma in proximity to the hamate bone and manifests as hypothenar hammer syndrome.

Auscultation

- Listen for a bruit in the supraclavicular fossa over the subclavian artery.
- When pulses are not palpable, a doppler is used to assess blood flow in the arteries.
- A blood pressure difference > 10 mmHg reflects hemodynamically significant stenosis in the innominate, subclavian, or axillary arteries. In these situations, the higher blood pressure is reflective of the patient's actual blood pressure.

Chest and Abdominal Exam:

Inspection

• The aorta usually is typically not visible on the exam. However, a large aneurysm may be seen pulsating between the xiphoid and umbilicus, especially in thin patients.

Palpation

- The aorta bifurcates at the level of the umbilicus. To palpate the aorta, press your fingers on both sides of the midline between the umbilicus and the xiphoid. To help relax the abdomen, ask the patient to bend their knees, flex their hips, and relax their abdominal muscle. The goal is not only to feel the aortic pulse but also to estimate the size of the aorta. In people without aortic aneurysms, a palpable aorta is often the size of the patient's thumb. A tender, enlarged, pulsatile abdominal mass may represent a symptomatic aortic aneurysm or inflammatory aneurysm.
- The sensitivity of palpation to detect an abdominal aortic aneurysm is low (29%) for small (3.0- 4cm) aneurysms. Moreover, even large aneurysms > 5 cm may not be detected on physical exam (sensitivity of 76%). False positives can be found in elderly patients who have tortuous anterior placed aorta. It is important to mention that palpation of an abdominal aortic aneurysm is safe and has never been reported to precipitate aortic rupture. When an aortic aneurysm is identified, A complete peripheral arterial examination should be performed looking for evidence of distal embolization, ischemia or associated peripheral artery aneurysms (femoral, popliteal).
- The iliac arteries lie deep in the pelvis and are usually not palpable, even if aneurysmal.

Auscultation

- Cardiac auscultation is performed to assess rate and rhythm with special attention to the presence of any arrhythmias, gallops, and murmurs.
- Bruits in the abdomen are associated with arterial stenosis. The origin of the bruit could be renal, mesenteric, or aortoiliac.

Lower Extremity Exam:

Inspection

- Pallor, cyanosis, and poor capillary refill are signs of chronic limb ischemia. Muscle atrophy, hair loss, and thick toenails may also be present.
- **Dependent rubor** and pallor with elevation indicate advanced peripheral occlusive disease. Dependent rubor is hyperemic erythematous discoloration of the limb in a dependent position (sitting or standing). That is, dependent rubor is due to maximally dilated capillaries and the effects of gravity. However, the limb becomes pale once the foot is elevated (the patient lies down). Dependent rubor is usually associated with rest pain and edema. It is frequently misdiagnosed as cellulitis.
- Ulcers need to be identified as neuropathic ulcers or ischemic ulcers. Neuropathic ulcers are at pressure points over the plantar aspect of the metatarsal head. Ischemic ulcers are more often towards the tip of the toes.
- Livedo Reticularis: Violaceous mottling of the skin with a reticular pattern of the skin of the arms and legs. The term "livedo racemosa" is used for cutaneous findings in inflammatory or thrombotic vascular disease patients.
- Acrocyanosis: is defined as bluish discoloration of the extremities due to high deoxygenated blood in the capillaries. It is a persistent disorder without episodic triphasic color response.
- Microembolic disease can manifest as blue to syndrome/trash foot.
- Dry skin is present in chronic limb ischemia because the sebaceous glands function poorly without adequate blood flow.
- Edema is called "pitting" when the indentation persists after applying pressure to a small area. *Pitting edema* is associated with systemic diseases like heart failure, chronic kidney disease, hypoproteinemia, or local disease of the veins or lymphatic. Non-pitting edema is observed when the indentation does not persist. It is associated with myxedema, lipedema, and advanced lymphedema.
- Claudication: Patients with claudication may have no significant finding on inspection apart from muscle atrophy or hair loss.

Auscultation

• Auscultate the femoral region for the presence of any bruits. Auscultation may also find continuous bruits which are likely due to an arterio-venous fistula.

Palpation

• Femoral pulse: palpated under the inguinal ligament, two-finger breaths from the pubic tubercle.

- Popliteal pulse: with the patient's knee flexed, both hands are wrapped around the knee, and the tips of the fingertips are pressed into the popliteal space (posteriorly). The pulse is located slightly lateral to midline. A normal popliteal artery may not be palpable.
- Dorsalis Pedis: palpated in the dorsum of the foot between the first and second extensor tendons. Located just lateral to the tendon of extensor hallucis longus. You can ask the patient to extend their great toe to identify the tendon. Postier tibial: palpated b posterior to the medial malleolus. It is easier to palpate with the foot passively dorsiflexed.
- Peroneal artery: not palpable. However, it can be found with a doppler on the lateral aspect of the ankle
- When a pulse is not palpated, a doppler is used to assess the blood flow. The doppler signal can be triphasic, biphasic, or monophasic.
- Triphasic and biphasic doppler signals indicate good blood flow.
- Monophasic signals correlate with a moderate to severe decrease in arterial blood flow.
- In severe ischemia, a soft continuous venous signal may be all that is heard.
- Temperature changes may help demarcate the level of disease.
- Sensory loss may be present in acute ischemia and chronic neuropathy.

Differential diagnosis of claudication

Table 1: **Table 1. Differential diagnosis of claudication** Adapted from: Dormandy JA, Rutherford RB. Management of peripheral arterial disease (PAD). TASC Working Group. TransAtlantic Inter-Society Concensus (TASC). J Vasc Surg 2000; 31:S1.

Condition	Location of pain or discomfort	Character of discomfort	Onset relative to exercise	Effect of rest	Effect of body position	Other features
Vascular claudica- tion	Muscles of the buttock, thigh, or calf. Rarely the foot	most common is cramping. May complain of aching, fatigue, weakness, or pain	onset of pain is after some degree of exercise	Relieved by rest.	None	Reproducible

Condition	Location of pain or discomfort	Character of discomfort	Onset relative to exercise	Effect of rest	Effect of body position	Other features
Nerve root compres- sion (eg, herniated disc)	Radiates down leg, usually posteriorly, usually from the back	Sharp lancinating pain. Electric.	Soon after, if not im- mediately after onset	Not quickly relieved. Often present at rest too.	Adjusting back position may relieve pain	History of back problems
Spinal stenosis	Hip, thigh, or buttock (within affected der- matome)	Motor weakness more prominent than pain	After standing for some length of time. Also may occur after walking confusing it with vascular claudication	Relieved by resting only if position changed	Relieved by lumbar spine flexion (sitting or stooping forward)	Frequent history of back problems, provoked by intraab- dominal pressure
Hip arthritis / knee arthritis	Hip, thigh, buttocks. Or knees	Aching discomfort, usually localized to hip and gluteal region or knees	After variable degree of exercise. Or standing.	Not quickly relieved (and may be present at rest)	More comfortable sitting (ie, weight taken off legs)	Variable, may relate to activity level, weather changes. Tenderness when pressing on hip or knee area
foot arthritic, inflamma- tory processes	Foot, arch	Aching pain	After variable degree of exercise	Not quickly relieved (and may be present at rest)	May be relieved by not bearing weight	Variable, may relate to activity level

Condition	Location of pain or discomfort	Character of discomfort	Onset relative to exercise	Effect of rest	Effect of body position	Other features
Venous claudica- tion	Entire leg, but usually worse in thigh and groin	Tight, bursting pain	After walking	Subsides slowly	Relief speeded by elevation of the extremity	History of iliofemoral deep vein thrombosis, signs of venous congestion, edema, venous stasis dermatitis or ulcers
Diabetic Neuropa- thy	numbness is sock like fashion in both feet	tingling, numbness.	constant	-	-	poorly controlled diabetes.
Night time cramps	foot, calf and thigh	crampy pain at night	not related to exercise	-	-	may be associated with use of diuretics.

Carotid Disorders

Carotid Artery Stenosis

Angela Kokkosis, MD, FACS and Michael Malinowski, MD, MEHP, FACS

Note

By the end of this chapter, students will:

- Review the foundational knowledge to appraise asymptomatic and symptomatic carotid artery stenosis.
- Identify and describe risk factors, symptoms, and signs of carotid artery disease.
- Review and describe indications and contraindications for medical and surgical management of carotid artery stenosis.
- Propose next best steps in patient work-up and treatment of carotid artery disease.
- Describe the evidence-based outcomes of surgical management of carotid artery disease.

Key Facts

- 1. Stroke is the leading cause of disability nationally and 3rd leading cause of death with a breakdown of 80% occlusive (ischemic) and 20% hemorrhagic.
- 2. Symptomatic carotid stenosis presents with only three symptoms including: stroke, transient ischemic attack (TIA) and amaurosis fugax (transient retinal ischemia).
- 3. Risk factors include: age, smoking, CAD, diabetes, hypertension, hyperlipidemia and genetic/family history.
- 4. Seminal studies include ACAS and NASCET. They define an 11% five-year risk of stroke for asymptomatic carotid artery stenosis (>60%) and 24% two-year risk of stroke for symptomatic disease (>70%), respectively.
- 5. The CREST landmark study associates carotid artery stenting with higher perioperative risk of stroke and carotid endarterectomy (CEA) with higher risk of myocardial infarction (MI) in symptomatic patients.
- 6. Carotid artery stenting (CAS) is indicated for recurrent stenosis after CEA, neck immobility, high carotid bifurcation, contralateral occlusion, high risk open surgical candidate due to cardiopulmonary comorbidity and neck radiation.

- 7. Transcarotid artery revascularization (TCAR) is available for direct common carotid delivery of stent while avoiding aortic arch manipulation and providing embolic protection through flow reversal.
- 8. There is a marginal benefit for intervention for asymptomatic carotid artery stenosis patients with significant cardiopulmonary disease due to risks of associated perioperative events.

Carotid Disease Slide Deck

Please find the slide deck corresponding to this eBook chapter here.

Please find a video recording of Dr. Kokkosis and Dr. Malinowski's lecture using the above slide deck here.

•

How We Suggest Using the Pre/Post Questions

The pre/post questions are listed below. They are all multiple choice questions with a single right answer. To best guide your learning, we have hidden the answers in a collapsible menu. Before reading the chapter, we suggest giving the questions a try, noting your answers on a notepad. After reading the chapter, return to the questions, reevaluate your answers, and then open the collapsible menu to read the correct answer and discussion. Do not fret if you have difficulty answering the questions before reading the chapter! By the end of the chapter, we are certain you will have covered the knowledge necessary to answer the questions. There will be a teaching case at the end of the chapter. This is another opportunity to exercise your new knowledge!

Pre/Post Questions

Case Based Questions

- 1. A 75-year-old male smoke presents with recent visual changes to his right eye that occurred yesterday. The patient reports a shading of his visual field that resulted in momentary monocular blindness followed by return to normal vision. He has a carotid duplex showing >50% diameter reduction to his right internal carotid artery and >80% stenosis to his left internal carotid artery. What treatment should be offered to his patient?
- A. Emergent DC cardioversion to treat any underlying arrythmia.
- B. Left carotid endarterectomy with shunt placement.
- C. Left carotid TCAR.
- D. Right carotid endarterectomy.

E. Placement on Apixaban and measurement of PF4 with medical management.

Answer

D. Right carotid endarterectomy

Discussion: The patient has evidence of amaurosis fugax with temporary monocular blindness as a sign of symptomatic right internal carotid artery disease. His left carotid artery is in a high-grade range and therefore is lower risk of stroke than the contralateral symptomatic lesion. There should be no immediate surgical treatment of the asymptomatic left carotid lesion. There is no evidence that the patient has an underlying arrythmia that needs cardioversion and medical management with Apixaban and PF4 levels are unrelated to this patient's current pathology.

- 2. A 50-year-old female patient with >80% right internal carotid artery stenosis presents to clinic for her first postoperative visit after carotid endarterectomy (CEA). She has no interval neurologic events since her discharge, has a soft neck with a clean incision. During your neurologic exam you notice an unintentional, subtle tongue deviation to the side of surgery. What is the most likely facial nerve involved in this finding?
- A. Vagus Nerve.
- B. Hypoglossal Nerve.
- C. Glossopharvngeal Nerve.
- D. Long Thoracic Nerve.
- E. Hering's Nerve.

Answer

B. Hypoglossal Nerve

Discussion: This patient has evidence of hypoglossal nerve neuropraxia which can occur as the result of injury or traction on the nerve during distal dissection of the internal carotid artery. Although both glossopharyngeal nerve injury and vagal injuries can also occur during CEA, the former causes oropharyngeal dysfunction with swallowing and the latter causes vocal cord paralysis leading to a hoarse voice. The long thoracic nerve is more related to thoracic outlet surgery and innervation of the serratus anterior. The Nerve of Hering is related to carotid sinus innervation and is unrelated to this clinical finding.

3. During the initial evaluation of a patient with high grade symptomatic carotid disease, you notice that the patient has internal carotid plaque on that side this is above the angle of the mandible at the 1st cervical vertebral body (C1). This appears to be too high to access through open surgery. The patient has no know history of coronary artery

disease, has a preserved ejection fraction and good functional status. He has a low-density lipoprotein level (LDL) of 200 mg/dL. What is the best treatment option for this individual?

- A. Transfemoral carotid artery angioplasty with placement on ASA only.
- B. Carotid artery enterectomy with shunt placement.
- C. Daily ASA therapy without any type of statin or antihypertensive therapy.
- D. Trancarotid artery revascularization (TCAR) surgery with dual antiplatelet therapy.
- E. No treatment is indicated.

Answer

D. Transcarotid artery revascularization surgery with dual antiplatelet therapy.

Discussion: This patient has symptomatic carotid stenosis with an overall optimal coronary health. Because his lesion is high at C1 and not surgically accessible, they would not qualify for a traditional carotid endarterectomy. Transfemoral carotid artery angioplasty is not indicated without stent placement, and daily ASA therapy without statin treatment for an LDL of 200mg/dL does not qualify as optimal medical management. Surgical treatment is indicated in this symptomatic patient since he has no significant cardiopulmonary disease and TCAR is the only option listed that could accomplish this outcome successfully.

- 4. A 60-year-old otherwise healthy woman, with no underlying comorbidities has a right carotid artery bruit on physical examination. She is concerned that she is at risk for stroke. She denies any episodes of vision changes, upper or lower extremity deficits, or speech impairments. She currently takes 81mg of aspirin daily, along with a multivitamin. What is the next step in management for this patient?
- A. No further treatment necessary.
- B. Carotid duplex.
- C. CT angiogram head and neck.
- D. Neurology evaluation.
- E. Addition of statin therapy.

Answer

A. No further treatment necessary.

Discussion: This patient has an incidental finding of a carotid bruit on physical exam,

however she has no risk factors for carotid disease (such as smoking, hyperlipidemia, smoking, family history, diabetes or hypertension). Additionally, she is neurologically asymptomatic. Therefore, carotid duplex, CTA, neurology evaluation, and the addition of statin therapy are not indicated. The prevalence of >75% carotid stenosis for those with a carotid bruit has been found to be very low at 1.2%.

- 5. A 65-year-old man who is right-handed is undergoing evaluation for a coronary artery bypass graft surgery (CABG). As part of his workup, a carotid duplex is performed which demonstrates a chronic right carotid occlusion and a >80% stenosis of his left carotid artery. He has no history of ocular or cerebrovascular events. He has hyper-cholesterolemia and well-controlled hypertension. What is the next best step for this patient?
- A. Proceed with the CABG as planned and continue medical therapy with aspirin and statin.
- B. Left carotid endarterectomy before the CABG.
- C. Place the patient on dual antiplatelet therapy, in addition to statin therapy.
- D. Left TCAR (transcarotid stent) after the CABG.
- E. Place the patient on anticoagulation.

Answer

B. Left carotid endarterectomy before CABG.

Discussion: It is standard of care to assess the carotid arteries prior to performing a CABG, with the goal of minimizing the risk of perioperative stroke. This patient has asymptomatic bilateral severe carotid disease (right occlusion and left >80% stenosis). Therefore, to reduce the risk of perioperative stroke, current guidelines recommend carotid revascularization prior or concomitant with the CABG. The patient does carry one high risk criterion for TCAR (coronary disease requiring revascularization), however this should not be performed after the CABG. Dual antiplatelet therapy is indicated in the event this patient undergoes carotid stenting, however medical management should not be the only management of his carotid disease. Lastly, anticoagulation has no role in atherosclerotic disease, such as carotid disease.

6. A 55-year-old woman has a past medical history of coronary artery disease status post coronary stenting in 2019, hypertension, hyperlipidemia, and previous smoking history of 60 pack-years. Her cardiologist sends her for a carotid duplex which demonstrates a 50-69% carotid stenosis on the left side, and mild atherosclerosis on the right side. She denies any prior signs or symptoms of stroke or transient ischemic attack. How should this patient's carotid disease be managed?

- A. Left carotid endarterectomy, along with aspirin/statin therapy.
- B. Left transfemoral carotid stent, along with dual antiplatelet/statin therapy.
- C. Aspirin and statin therapy **only**, and routine surveillance carotid duplexes.
- D. Aspirin, statin therapy, blood pressure management, and routine surveillance carotid duplexes.
- E. No further management is indicated.

D. Aspirin, statin therapy, blood pressure management, and routine surveillance carotid duplexes.

Discussion: This patient has asymptomatic 50-69% left carotid stenosis with the associated risk factors of vascular disease which include hypertension, hyperlipidemia, and smoking history. Medical therapy needs to be employed to reduce these factors, thus a single antiplatelet, statin, and blood pressure medication are indicated. Fortunately, the patient is not a current smoker, however if she was, then smoking cessation counseling would be added to her treatment plan. Carotid revascularization is indicated only for asymptomatic carotid stenosis that is >70% based on current guidelines.

Operative Footage Questions

These questions are associated with the carotid endarterectomy (CEA) footage (**short version**) found at the bottom of the chapter.

- 1. What is the first muscle layer encountered in a CEA (i.e. the first muscle deep to skin)?
- A. Sternocleidomastoid
- B. Digastric
- C. Platysma
- D. Scalene

Answer

C. Platysma

Discussion: The platysma is the most superficial muscle in the neck. It covers most of the anterior and lateral aspect of the neck. It is the first muscle layer encountered in the neck during a CEA. It will be bisected and repaired upon neck closure. While the sternocleidomastoid is superficial, it is deep to the platysma. The sternocleidomastoid is an important landmark as it forms the anterolateral boundary of the carotid triangle.

You will dissect along its medial border and retract it laterally in order to access the carotid sheath. The digastric muscle is a small, "two-bellied" muscle located under the mandible. The posterior belly forms the superior border of the carotid triangle. It is often visualized in patients with high carotid bifurcations. The scalene muscles are deep to the sternocleidomastoid muscles and lateral to the cervical spine. They are not manipulated in a CEA.

- 2. What structure is **not** found in the carotid sheath?
- A. Internal jugular vein
- B. External jugular vein
- C. Common carotid artery
- D. Vagus nerve

Answer

B. External jugular vein

Discussion: The external jugular vein is not in the carotid sheath. It runs superficial to and obliquely across the sternocleidomastoid before passing to the posterior border of the sternocleidomastoid as it descends deep into the neck. The carotid sheath contains the common carotid artery anteromedially, the internal jugular vein anterolaterally, and the vagus nerve posteriorly.

- 3. In what order do you **unclamp** the carotid vessels at the end of a carotid endarterectomy?
- A. External carotid artery \rightarrow common carotid artery \rightarrow internal carotid artery
- B. Common carotid artery \rightarrow internal carotid artery \rightarrow external carotid artery
- C. Internal carotid artery \rightarrow external carotid artery \rightarrow common carotid artery
- D. Internal carotid artery \rightarrow common carotid artery \rightarrow external carotid artery

Answer

D. External carotid artery \rightarrow common carotid artery \rightarrow internal carotid artery

Discussion: In conventional CEA, the internal carotid artery (ICA), the common carotid artery (CCA), and the external carotid artery (ECA) are clamped in that order such that any atherosclerotic debris loosened during vessel manipulation and/or clamping embolizes into the ECA where it could cause extracranial ischemia rather than the ICA where it could cause intracranial ischemia (i.e. ischemic stroke). A useful mnemonic to remember

the clamping order is **ICE** for **ICA**, **CCA**, and **ECA**. The order when unclamping is the opposite, **ECI**. The rational is similar. By unclamping the ECA first, remaining debris or debris loosened during removal of the clamps will flow up the ECA rather than the ICA. By unclamping the CCA second, and dislodged debris will flow down the open ECA rather than the ICA. The ICA is unclamped last. Options A and B are dangerous as they do not minimize the risk of embolus flowing into the ICA. Option C is simply incorrect.

Introduction

Hemispheric stroke related to carotid artery stenosis is a leading cause of both disability and death in the United States. Underlying etiologies for stroke include occlusive or hemorrhagic events with roughly 80% being related to occlusive pathology through embolus or in-situ thrombosis, the remaining 20% attributable to hemorrhage. Roughly 15% of stroke victims have a transient ischemic attack (TIA) that fully resolved prior to a later stroke event. Risk factors for carotid plaque formation are related to age, smoking, coronary artery disease, diabetes, hyperlipidemia, hypertension and family history of stroke. Due to carotid bulb anatomy, the most common area of plaque formation is within the proximal internal carotid artery. As plaque stenosis increases over time, the systolic velocity increases to maintain flow volumes which intensifies shear stress. This shear stress increases likelihood of plaque rupture, platelet aggregation and thromboembolization. There are multiple seminal studies that describe cohort comparisons of asymptomatic and symptomatic carotid artery stenosis with outcomes related to optimal medical management alone or as adjunct to surgical repair. Symptomatic carotid stenosis is described as carotid stenosis >50% with unilateral stroke, TIA or amaurosis fugax on the side of carotid disease. Amaurosis fugax is historically described as shade coming down across one eye on the side of stenosis to produce partial or complete, painless monocular visual loss related to transient retinal ischemia. Amaurosis fugax may be bilateral in the case of bilateral, symptomatic carotid artery stenosis.

Etiology

Atherosclerosis is the most common cause for the development of carotid artery disease. This process is defined by deposition of lipid-laden plaque at the carotid bifurcation, and potentially across a larger territory of the common carotid, external carotid, and internal carotid arteries. This plaque may contain varying degrees of calcification and/or thrombus. The mechanisms

by which atherosclerosis at the carotid bifurcation may lead to stroke or TIA are: occlusion (cessation of blood flow to the internal carotid artery) or embolization (plaque debris break off and travel through the internal carotid artery to the brain). There are various risk factors which may contribute to the degree of atherosclerosis and its progression. These include history of cigarette smoking, hyperlipidemia, coronary artery disease, diabetes, hypertension, advanced age, and family history of carotid disease or stroke.

Diagnostics and Imaging

Three primary imaging modalities are used to evaluate carotid artery stenosis with the lowest cost option being color flow duplex ultrasound (DSA) that allows a physician to determine peak systolic and end diastolic velocities throughout the carotid bifurcation. Based on the internally validated vascular laboratory criteria of the institution, these velocities can be correlated to ranges of degree of stenosis, with high grade stenosis defined as >70-80%. Since the modality is based primarily on velocity range, it cannot give exact stenosis such as 66%. The modality can also provide adjunct information about the blood flow waveforms in each arterial segment, as well as whether that flow is laminar or turbulent with utilization of color flow imaging. Limitations of this imaging include technician skill, inability to obtain optimal angle of Doppler interrogation for velocity determination, shadowing from heavily calcified lesions, poor visualization due to patient habitus and tortuosity.

Axial imaging options include both computed tomography angiogram (CTA) and magnetic resonance angiogram (MRA). Both of these options require some form of intra-arterial contrast, either iodinated contrast or gadolinium, respectively. However, they offer a fuller perspective of relevant anatomy and a more precise determination of stenosis within the limitations of the modality, with MRA often overestimating the degree of stenosis due to intrinsic properties of MRA imaging acquisition. Both CTA and MRA, although superior to DSA in determination of exact degree of plaque stenosis and arterial anatomy, sacrifice the physiologic information offered through DSA that speak to flow patterns, flow direction and turbulence. Axial imaging of CTA and MRA can define patency but do not speak to the dynamic nature of blood flow or directionality of flow.

Definitive determination of flow and directionality can be augmented to a carotid artery stenosis workup by diagnostic angiography. This requires femoral artery access and includes contrast administration, as well as a small risk of periprocedural embolization. However, it offers additional physiologic evaluation that might not be present in DSA and that is inherently lacking in CTA and MRA studies.

Asymptomatic Carotid Atery Stenosis Screening

The 2021 SVS clinical practice guidelines outline the following recommendations. In asymptomatic patients who qualify for carotid artery stenosis screening, duplex ultrasound is the recommended choice over CTA, MRA, or other imaging modalities.

Routine screening is **not recommended** for clinically asymptomatic carotid artery stenosis for individuals **without significant risk factors** for carotid disease.

Screening is **recommended** for clinically **asymptomatic carotid artery stenosis in in-dividuals** with significant risk factors** for carotid disease. High-risk groups include:

- Patients with lower extremity peripheral artery disease (PAD)
- Patients undergoing coronary artery bypass surgery (CABG)
- Patients aged >= 55 years with at least two traditional atherosclerotic risk factors (hyperlipidemia, hypertension, etc.)
- Patients aged >= 55 years and active cigarette smoking
- Patients with diabetes, hypertension, or coronary artery disease (CAD)
- Patients with clinically occult cerebral infarction noted on brain imaging studies.

The presence of a carotid bruit increases the likelihood of detecting significant stenosis. Asymptomatic patients with an abdominal aortic aneurysm (AAA) or previous radiotherapy to the neck who do not meet the criteria of any of the high-risk groups above do not require screening. It has been shown that the prevalence of carotid stenosis increases proportionally with the number of risk factors present.

Treatment

Carotid Artery Endarterectomy (CEA)

This procedure has been performed since the 1950s, either by plaque endarterectomy and patch angioplasty or primary arterial repair. To prevent arterial restenosis, patch angioplasty has become the standard of arterial closure after plaque removal. The procedure can involve cerebral monitoring including electroencephalography (EEG), transcranial Doppler (TCD) and stump pressure monitoring or be performed awake to directly monitor patient motor response. Endarterectomy and patch repair can be performed under a "clamp and sew" mentality or with an arterial shunt to maintain cerebral perfusion. Risks include cardiopulmonary risk of acute myocardial ischemia, <3% perioperative risk of neurologic event, neck hematoma or cranial nerve injury of roughly 5-10% affecting the vagus, marginal mandibular, recurrent laryngeal or hypoglossal nerves.

Carotid Artery Stenting (CAS)

Transfemoral Carotid Artery Stenting (TFCAS) with Embolic Protection

Transfemoral stenting requires some type of protection from embolization including a distal internal carotid artery retrievable filter or flow arrest procedure to prevent cerebral embolization during stent placement with or without angioplasty.

Transcarotid Artery Revascularization (TCAR)

Treatment of a carotid stenosis that avoids aortic arch manipulation involving direct common carotid artery exposure and sheath placement to allow for transcarotid stent delivery to the internal carotid artery. The common carotid artery sheath is connected to a femoral vein sheath so that the natural arterial pressure gradient reverses flow across the distal internal carotid artery driving blood and possible embolus into the arterial tubing circuit and across a filter before it reenters the venous circulation. As FDA approval for the device was delivered in 2016, the technology is less than 10 years old without robust long-term follow up data. This technique offers a lower perioperative stroke risk than transfemoral stenting, for multiple reasons including lack of transaortic arch manipulation and great vessel cannulation which can result in embolus prior to placement of an internal carotid artery embolic protection device.

Optimal Medical Management

Understandably, optimal medical management requires full risk evaluation of the individual patient in question including other comorbidities, drug allergies, compliance, etc. We have listed a few broad recommendations to follow that offer general guidance surrounding the dynamic target of optimal medical management for arterial disease.

Antiplatelet Therapy

- ASA offers a 22% risk reduction in major vascular events with no difference in protection based on dosage (81 versus 325 mg).
- Clopidogrel can be used as an adjunct or alternative to ASA, but the added benefit from dual antiplatelet combination in asymptomatic carotid artery stenosis is unproven.

Anticoagulants

• Only useful for prevention of cardioembolic strokes due to arrhythmia or prosthetic valve.

Hypertension Treatment

• Recommended blood pressure range of <130/80 with individual antihypertensive regimen based on other comorbidities and patient risk factors.

Diabetic Control

• In accordance with best practice for diabetes management, the patient's hemoglobin A1c should be <7.0

Smoking Cessation

• Treatments offered include nicotine replacement therapy (NRT), varenicline or bupropion as first line agents.

Hyperlipidemia Management

• Regimen goals of LDL <100mg/dl, or <70mg/dl depending on risk profile.

Outcomes and Surveillance

Asymptomatic carotid stenosis

- The historically touted Asymptomatic Carotid Atherosclerosis Study (ACAS) demonstrated that patients with >60% carotid stenosis who underwent CEA benefited significantly from stroke risk reduction at 5 years (5.1% for CEA vs. 11% for optimal medical therapy consisting of aspirin alone).
- More recent studies suggest that with the current optimal medical management, which consists of antiplatelet medication and statin therapy, 5-year stroke risk is highest in patients with >70% carotid stenosis, and therefore this patient population would benefit from carotid endarterectomy.
- Patients who are deemed high risk, either due to an anatomic (such as surgically inaccessible bifurcation or restenosis after previous CEA) or physiologic findings (congestive heart failure, severe coronary artery disease, or chronic obstructive pulmonary disease), may be considered for TCAR given the equivocal results of perioperative stroke or death at 1.3%, as compared to CEA.

• Asymptomatic patients with significant risk factors found to have moderate stenoses (50%-79%) should be followed every 6-months to detect disease progression. High risk patients with <50% stenosis can be followed-up annually.

Symptomatic carotid stenosis

- Patients who have >50% carotid stenosis and have developed symptoms of TIA or stroke were found to benefit from CEA in the pivotal North American Symptomatic Carotid Endarterectomy Trial (NASCET) because of the significant 2-year stroke risk reduction as compared to optimal medical management (15.7% vs. 22.2%.). An even greater stroke risk reduction was seen in patients with >70% carotid stenosis (9% CEA vs. 26% medical management).
- Current management of patients with symptomatic >50% carotid stenosis who are low/standard risk is carotid endarterectomy over transferoral carotid stenting (TFCAS), as there are no studies to date which have shown benefit of TFCAS.
- Patients who are deemed high risk, as defined above, may be considered for TCAR over TFCAS due to the significantly lower incidence of in-hospital stroke and death (1.6% vs. 3.1%).
- Post-operative surveillance (by duplex ultrasound) after open (CEA) or endovascular (TF-CAS) repair of the carotid artery is strongly recommended by the SVS to monitor for signs of restenosis in the repaired artery or atherosclerotic disease progression in the unoperated, contralateral artery. Duplex ultrasound testing is recommended within 30 days of the procedure, then every 6 months for 2 years, then annually.
- Restenosis <50% warrants the regular surveillance protocol; 50-99% warrants closer follow-up, confirmation with a CTA, and possible angiographic evaluation; 100% restenosis warrants surveillance and medical treatment of the contralateral carotid artery.
- It should be noted, there is some debate as to the economic and medical value of continuing post-operative duplex ultrasound surveillance after successful CEA with patch closure when the immediate post-operative duplex was normal or showed minimal disease.

Teaching Case

Scenario

An 81 year old male with a significant smoking history and prior three vessel CABG five years ago, presents with monocular right eye blindness that occurred two days ago. He has no prior ophthalmologic conditions and states that he describes the process of a veil coming down over

his right eye with resolution about a minute later with complete return of normal vision at that point. He denies any other symptoms during the event or since, such as motor or sensory deficits, speech, etc. He did not think much of the event but presented after his wife told him to see someone about the event.

Exam

HEENT: No prior neck incisions, good cervical extension.

Cardiac: Regular rate and rhythm. Healed sternotomy scar.

Pulmonary: Clear to auscultation throughout.

Abdominal: Soft and nontender.

Neurologic: All cranial nerves 2-12 intact, no lateralizing deficits, 5/5 strength to all extrem-

ities.

Optho: No visual deficits at 20 feet from eye chart.

Imaging

Duplex Ultrasound (Peak Systolic Velocity/End Diastolic Velocity)

Location	Right	Left
Proximal ICA	540/240 cm/s	120/45 cm/s
Mid ICA	230/145 cm/s	119/37 cm/s
Distal ICA	240/110 cm/s	110/23 cm/s

Duplex Report: Based on color flow duplex imaging there is evidence of 80-99% stenosis of the right internal carotid artery segment and <50% stenosis to the contralateral side.

Discussion Points

N.B. There is no prepared answers for the questions below.

However, we feel this chapter contains all the necessary information to answer the questions. If not, please let us know!

1. Please explain the pathophysiology of the visual event for this patient? Describe why it can be termed amaurosis fugax. Ensure understanding that amaurosis fugax is the result of carotid plaque embolization to the retina.

- 2. Please list the patient's risk factors for carotid disease? What is best medical management to optimize these risk factors?
- 3. Is this patient asymptomatic or symptomatic based on the clinical scenario presented?
- 4. What next steps should be pursued to offer effective and timely treatment to this patient? Please discuss adjunct imaging such as CTA or MRA to determine anatomic characteristics of the lesions such as ulceration, vessel patency, level (accessible or high lesions), etc.
- 5. What surgical managements could be suggested to this patient? Please include a discussion of carotid endarterectomy, transfemoral stenting or TCAR.
- 6. What medications should be started in this scenario? Please consider ASA, Plavix, statin medications, etc.
- 7. What are some possible relevant complications of surgical intervention, including periprocedural stroke risk?

Key Articles

- 1. Ricotta JJ, Aburahma A, Ascher E, Eskandari M, Faries P, Lal BK; Society for Vascular Surgery. Updated Society for Vascular Surgery guidelines for management of extracranial carotid disease. J Vasc Surg. 2011 Sep;54(3):e1-31.(Ricotta et al. 2011)
- 2. AbuRahma AF, Avgerinos EM, Chang RW, Darling RC 3rd, Duncan AA, Forbes TL, Malas MB, Murad MH, Perler BA, Powell RJ, Rockman CB, Zhou W. SOCIETY FOR VASCULAR SURGERY CLINICAL PRACTICE GUIDELINES FOR MANAGEMENT OF EXTRACRANIAL CEREBROVASCULAR DISEASE. J Vasc Surg. 2021 Jun 18. (AbuRahma et al. 2022)
- 3. Endarterectomy for asymptomatic carotid stenosis. Executive Committee for the Asymptomatic Carotid Atherosclerosis Study. JAMA 1995;273(18):1421-8. (Walker 1995)
- 4. Beneficial effect of carotid endarterectomy in symptomatic patients with high-grade stenosis. North American Symptomatic Carotid Endarterectomy (NASCET) Trial Collaborators. N Engl J Med 1991;325(7):445-53. ("Beneficial Effect of Carotid Endarterectomy in Symptomatic Patients with High-Grade Carotid Stenosis" 1991)
- 5. Howard D.P.J., Gaziano L., Rothwell P.M.: Risk of stroke in relation to degree of asymptomatic carotid stenosis: a population-based cohort study, systematic review, and meta-analysis. Lancet Neurol 2021; 20: pp. 193-202.(Howard, Gaziano, and Rothwell 2021)

56

Additional Resources

Audible Bleeding Content

- Audible Bleeding Exam Prep: Cerebrovascular Chapter
- Audible Bleeding has an episode covering the NASCET trial. Listen to it below and find additional information here, or find the episode wherever you listen to podcasts.
- The Audible Bleeding Holding Pressure Series has an episode about carotid endarterectomy. The Holding Pressure Series is designed specifically for medical students! Listen to the episode below and find additional information here, or find the episode wherever you listen to podcasts.

	bs	

• TeachMe Surgery: Carotid Artery Disease

Serious Games

Touch Surgery Simulations.

- Must download the Medtronic Touch Surgery mobile application to access the modules.
 Available for Apple and Android mobile devices.
- Carotid Endarterectomy
- Carotid Artery Stenting

Gore Combat Manual

The Gore Medical Vascular and Endovascular Surgery Combat Manual is an informative and entertaining read intended as a vascular surgery crash course for medical students, residents, and fellows alike. Highly accessible with a thoughtfully determined level of detail, but lacking in learning activities (e.g. questions, videos, etc.), this resource is a wonderful complement to the APDVS eBook.



Please see pages 119-125.

Operative Footage

Developed by the Debakey Institute for Cardiovascular Education & Training at Houston Methodist. *YouTube account required as video content is age-restricted.* Please create and/or log in to your YouTube account to have access to the videos.

Carotid Endarterectomy (Short Version)

Carotid Endarterectomy (Long Version) Part 1

Carotid Endarterectomy (Long Version) Part 2

Carotid Endarterectomy (Long Version) Part 2

Transcarotid Artery Revascularization (TCAR)

Aortic Disorders

Abdominal Aortic Aneurysm (AAA)

Erin K. Greenleaf, MD, MS, Jonathan Bath, MD, Dawn M. Coleman, MD, FACS and Ezra Schwartz, MD, CM, MS.

Note

By the end of this chapter, students will:

- Review abdominal agrtic anatomy and aneurysm pathophysiology.
- Recognize history, physical exam, and radiologic findings typical of patients presenting with an abdominal aortic aneurysm.
- Identify indications for intervention on abdominal aortic aneurysms and the evidence-based foundation for these recommendations.
- Describe the basic procedural steps of both an endovascular and an open surgical approach for abdominal aortic aneurysm repair.
- Delineate appropriate surveillance intervals for imaging an abdominal aortic aneurysm, both before and after intervention.

Key Facts

- 1. Abdominal aortic aneurysms occur in 2-8% of the population in Western countries, with a slightly higher prevalence among males relative to females.
- 2. The average size of an abdominal aorta is approximately 2cm; hence, an aneurysm is diagnosed when the diameter of the aorta in this segment reaches 3cm (i.e. 1.5x normal diameter).
- 3. Risk factors for an abdominal aortic aneurysm growth include age, male sex, white race, smoking, atherosclerosis, family history of arterial aneurysms, and personal history of arterial aneurysms.
- 4. According to vascular surgery guidelines, elective repair of a fusiform abdominal aortic aneurysm is recommended once the aneurysm has reached 5.5cm diameter in males and 5 cm diameter in females, measured from outer wall to outer wall. Circumstances in which repair is indicated for smaller aneurysms include those associated with symptoms and/or rupture, saccular morphology, rapid ex-

- pansion, or those in females (as mentioned above), immunosuppressed patients (i.e. chemotherapy, radiation therapy, or solid organ transplantation) and patients with aortopathies (e.g. Marfan Syndrome, Ehlers-Danlos Syndrome, etc.).
- 5. Nearly 80% of abdominal aortic aneurysms are currently repaired via an endovascular approach, with an open surgical approach generally reserved for patients in which a stent graft would perform poorly.
- 6. Outcomes following endovascular abdominal aortic aneurysm repair have improved over time, as experience has accrued with endovascular technique. This pattern of progressive improvement has not similarly been seen for open surgical repair.
- 7. Surveillance regimens for patients with unrepaired abdominal aortic aneurysms suggest imaging via ultrasound or computerized tomography (CT) scan every three years if the aneurysm is 3.0-3.9cm diameter, every year if the aneurysm is 4.0-4.9cm, and every six months if the aneurysm is 5.0-5.4cm.

• How We Suggest Using the Pre/Post Questions

The pre/post questions are listed below. They are all multiple choice questions with a single right answer. To best guide your learning, we have hidden the answers in a collapsible menu. Before reading the chapter, we suggest giving the questions a try, noting your answers on a notepad. After reading the chapter, return to the questions, reevaluate your answers, and then open the collapsible menu to read the correct answer and discussion. Do not fret if you have difficulty answering the questions before reading the chapter! By the end of the chapter, we are certain you will have covered the knowledge necessary to answer the questions. There will be a teaching case at the end of the chapter. This is another opportunity to exercise your new knowledge!

Pre/Post Questions

Case Based Questions

- 1. A 75-year-old male with history of hypertension, hyperlipidemia, diabetes and degenerative disk disease presents to the emergency department with abdominal pain and a blood pressure of 90/60 mm Hg. The emergency department physician already obtained a CT scan showing what appears to be a contained rupture of an infrarenal abdominal aortic aneurysm. What is the next step in management?
- A. Resuscitate the patient to ensure a systolic blood pressure of at least 120 mm Hg.
- B. Two large bore IVs and admission to the ICU for closer monitoring.
- C. Transfer to the operating room for emergent repair of the aneurysm.

- D. Corroboration of the CT scan with an abdominal ultrasound.
- E. Contact the next of kin to request their presence at bedside for a family meeting.

C. Transfer to the operating room for emergent repair of the aneurysm.

Discussion: Explanations to come shortly

- 2. An 83-year-old female presents to your clinic with a 5.1cm abdominal aortic aneurysm. She has a history of chronic tobacco abuse, hyperlipidemia, osteoarthritis, and renal cell carcinoma status post left nephrectomy. She denies abdominal pain and is without significant physical exam findings. Based on her presentation, what is your recommendation to her?
- A. Continue surveillance with imaging every 6 months until her aneurysm reaches 5.5cm in diameter.
- B. Discuss repair of the aneurysm at this size.
- C. Prescribe an antiplatelet, statin, and beta blocker prior to discussing operative repair of the aortic aneurysm.
- D. Continue surveillance with imaging every 12 months until her aneurysm reaches 5.5cm in diameter.
- E. Tobacco cessation should be confirmed prior to committing to aneurysm repair.

Answer

B. Discuss repair of the aneurysm at this size.

Discussion: Explanations to come shortly

- 3. A 68-year-old male with history of hypertension, hyperlipidemia, tobacco abuse, obesity, and gout is found to have a 5.9cm infrarenal abdominal aortic aneurysm with a 21mm neck, patent common femoral arteries, but highly calcified/nearly occluded bilateral iliac arteries. Upon discussing repair with him in your surgery clinic, what would you advise the patient about his repair?
- A. He should undergo endovascular repair because there is at least 20mm of neck below the lowest renal artery before the aneurysm begins.
- B. He should undergo endovascular repair with percutaneous access because his common femoral arteries are widely patent.
- C. He should undergo endovascular repair because this will minimize his inpatient stay while maximizing the primary patency of the stent graft.

- D. He should undergo open repair because his diseased iliac arteries will preclude safe passage of sheaths and a device to the level of the aorta.
- E. He should undergo open repair because this will minimize his short-term morbidity.

D. He should undergo open repair because his diseased iliac arteries will preclude safe passage of sheaths and a device to the level of the aorta.

Discussion: Explanations to come shortly

- 4. A 78-year-old female is referred by her primary care physician after she was incidentally found to have a saccular aneurysm immediately distal to her right renal artery, which is the lower of the bilateral renal arteries. When documenting the location of this aneurysm, what terminology should be used to label it?
- A. Infrarenal
- B. Juxtarenal
- C. Suprarenal
- D. Infrainguinal
- E. No specific term exists

Answer

B. Juxtarenal

Discussion: Explanations to come shortly

- 5. Two days following an open transperitoneal repair of an infrarenal abdominal aortic aneurysm, an 81-year-old male with a history of a right colectomy for colon cancer has bright red blood in his stool. He remains in the ICU with a leukocytosis and appropriately tender abdomen without peritonitis. What is the next best step in management?
- A. Broad spectrum antibiotics and flexible sigmoidoscopy.
- B. Broad spectrum antibiotics and CTA.
- C. Observation as this is likely related to his previous colon resection.
- D. NPO order and immediate return to the operating room.
- E. Stool softeners to mitigate the risk of hemorrhoids and anal fissures.

A. Broad spectrum antibiotics and flexible sigmoidoscopy

Discussion: Explanations to come shortly

- 6. A 70-year-old female is undergoing an endovascular repair of an abdominal aortic aneurysm in the operating room under general anesthesia and a completion aortogram was just taken. A late blush is noted in the sac, seemingly arising from a lumbar artery. What kind of endoleak would this suggest is present?
- A. Type Ia endoleak; blood is filling the sac from the proximal extent of the stent graft.
- B. Type III endoleak; blood is filling the sac from the middle and can appear to be coming from a lumbar artery.
- C. Type Ib endoleak; blood is filling the sac retrograde from the iliac arteries and so is only seen with a late aortogram view.
- D. Type II endoleak; flow from a lumbar artery would cause a late blush and can be followed conservatively provided that the sac does not continue to expand.
- E. No endoleak.

Answer

D. Type II endoleak; flow from a lumbar artery would cause a late blush and can be followed conservatively provided that the sac does not continue to expand.

Discussion: Explanations to come shortly

Operative Footage Questions

10 come snortly.	Stay tuned:		

Introduction

The aorta is invariably the largest arterial conduit in the body, serving as the channel through which blood from the heart travels to every other body part. The wall of the aorta is comprised of three layers, which include the intima, media and adventitia. Its typical course begins at the distal side of the aortic valve and terminates at the aortic bifurcation, where it splits into

the two common iliac arteries. The thoracic aortic segment provides blood flow to the great vessels of the aortic arch, ensuring perfusion to the head and bilateral upper extremities via the innominate artery, the left carotid artery, and the left subclavian artery (in that order). The abdominal segment is a retroperitoneal structure that enters the abdominal cavity through the aortic hiatus of the diaphragm, at the level of T12. The branches of the abdominal aorta include, in descending order, the two inferior phrenic arteries, the celiac axis or trunk, the superior mesenteric artery (SMA), the renal arteries, the inferior mesenteric artery (IMA), and ultimately, the bilateral common iliac arteries. The branches of both the thoracic and abdominal aorta provide blood flow to areas of critically vital viscera.




Figure 1: Anatomically normal agrta with main and visceral branches.

While the aorta serves a critical role for the rest of the body, it can be the site of significant pathology. Aneurysms, which are common pathology in both the thoracic and abdominal aorta, are defined by an increase in arterial diameter to 1.5 times the normal diameter of the vessel. As

the diameter of the aortic aneurysm increases, the risk of rupture increases in direct proportion. For example, a patient with a 4cm abdominal aortic aneurysm (AAA) has an approximate 1-year rupture risk of 1%; a patient with a 5.5cm aneurysm has up to a 10% 1-year rupture risk. With the widespread use of cross-sectional imaging, many aortic aneurysms are discovered incidentally. Others are found on screening exams. The US Preventative Services Task Force recommends that men aged 65 to 75 years of age who have ever smoked tobacco undergo an abdominal ultrasound to screen for aneurysmal disease. Aortic aneurysms may also present symptomatically, with chest, abdominal and/or back pain and occasionally a palpable pulsatile abdominal mass, which is widely believed to represent an increasingly unstable aneurysm.

Etiology

Certain risk factors, some modifiable and others non-modifiable (a.k.a. innate), predispose individuals to the development of aortic aneurysms. These include age, male gender, family history of aneurysms or aortopathies, and comorbidities including hypertension, hyperlipidemia, and peripheral vascular disease. Interestingly, female sex, African American race, and a history of diabetes have been found to be protective against the development and rupture of aortic aneurysms. A personal history of Marfan's syndrome, Loeys Dietz syndrome, or Ehlers Danlos Type IV (a.k.a. vascular Ehlers-Danlos or vEDS) is associated with a particularly elevated risk of aneurysmal degeneration. Medical providers should counsel and assist their patients to mitigate all potentially modifiable risk factors.

Diagnostics and Imaging

When evaluating a patient with a suspected aortic aneurysm, a thorough history, physical examination, and imaging will help to narrow the differential diagnosis. Most often, patients present without symptoms; however, if symptomatic, typical complaints include chest, abdominal or back pain. Physical exam findings may not be overt but vital signs may provide a first clue in a symptomatic patient as heart rate and blood pressure might deviate from normal ranges. Patients with thoracic aortic aneurysms (TAA) often have no other easily identifiable physical exam findings aside from vital sign derangements in those with symptomatic or ruptured aneurysms. In thin patients with suspected AAA, the abdominal exam may demonstrate a pulsatile mass at or adjacent to the midline.





Figure 2: Aneurysm Topography

Although the history and physical exam are indispensable to patient evaluation, the diagnosis of an aneurysm is usually secured with imaging. The mainstay of imaging is computed to-mography (CT), while ultrasound, magnetic resonance imaging (MRI), and angiography are useful modalities in non-urgent or non-emergent settings. The benefits of CT, particularly the richness of anatomic data obtained, must be weighed against the potential risks related to contrast administration and exposure to radiation. Since the 2019 statement from the U.S. Preventive Services Task Force regarding one-time screening with ultrasonography in men aged 65 to 75 years who have ever smoked, the use of ultrasound has a more prominent role in AAA screening. Equipped with history, exam, and imaging data, the diagnosis of an aortic aneurysm can usually be made.

When looking at an imaging study, abdominal aortic aneurysms are labeled in regard to their location relative to the renal arteries:

- Aortic aneurysms are termed "suprarenal" when their most proximal extent is above the level of the renal arteries. Suprarenal abdominal aortic aneurysms typically involve visceral branches of the aorta as well (i.e. the SMA or celiac trunk) and thus may also be classified as a Crawford type IV thoracoabdominal aortic aneurysm (TAAA)
- Aneurysms are labeled "pararenal" when the proximal extent of the aneurysm is at the level of the renal arteries and includes the origins of the renal arteries.

- Abdominal aortic aneurysms are termed "juxtarenal" when their most proximal extent is immediately below the take-off of the renal arteries.
- Lastly, aneurysms are labeled "infrarenal" when they begin below the renal arteries, often with a neck of non-dilated aorta between the lowest renal artery and the top of the aneurysm. As one might expect, the location of the aneurysm has implications for treatment options.



AAA anatomy description relative to the renal arteries.



Please see Figure 2 of this American Heart Association publication depicting the anatomic classifications of AAAs.

Treatment

Once a diagnosis has been obtained, considerations for management follow. Similar to a thoracic aortic aneurysm (TAA), indications for repair are reflective of risk for rupture. The aim is to intervene when the rupture risk exceeds the risks posed by surgery. Intervention

is considered when an aneurysm reaches the size threshold of 5.5cm in males and 5cm in females or if the growth rate exceeds 5 mm in 6 months. Again, this threshold is modified in patients with a strong family history of aneurysmal disease or collagen vascular disorder, or when patients present symptomatically or with a ruptured aorta. The topography of the aneurysm must also be considered as a saccular aneurysm, an isolated pouch off the sidewall of the aorta, presents a much greater risk of rupture and should be fixed when identified.

On first evaluation, patients with AAAs should be medically optimized to minimize their perioperative risk of morbidity and mortality. Multiple factors must be considered when deciding between an open surgical approach and an endovascular one. While the prevalence of endovascular aortic repair (EVAR) has exponentially increased in the past two decades relative to open surgical repair (OSR), each aneurysm poses a unique set of potential constraints that suggest a preference for one type of repair over another.

Please see the Section section and watch the included videos.

Endovascular Aortic Repair (EVAR)

Endovascular aortic repair (EVAR) is a novel approach to AAA treatment that has gained widespread popularity since its first successful use by Dr. Juan Parodi and Dr. Julio Palmaz in 1991. EVAR necessitates cross-sectional imaging to measure the AAA dimensions and determine the appropriate size of an aortic stent graft. Stent grafts are straight or bifurcated tubes with a wire framework covered by non-porous fabric. Simply, it is a tube that can be deployed from inside a vessel.

EVAR Procedural Steps

A step-by-step guide of a typical, uncomplicated EVAR for an infrarenal AAA is listed below. Please see the Section section to a video guide of a typical EVAR.

- 1. Bilateral femoral access via percutaneous or open surgical exposure (a.k.a. "femoral cut-downs")
- 2. Heparinization to achieve an activated clotting time (ACT) greater than 250 seconds.
- 3. Placement of stiff wires into the abdominal agree and upsizing of sheaths into the access vessels.
- 4. Aortogram to visualize renal arteries and deployment of the endograft main body under the origins of the renal arteries.
- 5. Cannulation of contralateral limb gate using a wire and catheter with subsequent confirmation that both the wire and catheter are within the main body.

- 6. Aortoiliac angiogram to measure the distance from the contralateral gate to the contralateral hypogastric (i.e. internal iliac) artery.
- 7. Deployment of contralateral limb with the distal landing site just proximal to the hypogastric artery.
- 8. Partial deployment of the ipsilateral limb.
- 9. Aortoiliac angiogram to measure the distance from the ipsilateral gate of the main body to the ipsilateral hypogastric artery.
- 10. Deployment of the remainder of the ipsilateral limb.
- 11. Ballooning of the proximal and distal landing zones, and areas of stent overlap.
- 12. Remove wires and sheaths and close arteriotomies (often with closure devices).
- 13. Reverse heparin with protamine.
- 14. Apply pressure to surgical sites and monitor for hematoma formation.

EVAR Considerations and Planning


The need for imaging can be a limiting factor such as when patients with a ruptured aneurysm are too hemodynamically unstable to stop at the CT scanner prior to repair or when a patient's kidney function is too impaired to tolerate a large contrast bolus.

If the situation is amenable to getting a CT angiogram to measure anatomic dimensions, several components of the aneurysm anatomy should be assessed to determine anatomic suitability for EVAR. These include the length, width, angulation of the aneurysm, and presence of disease at the aneurysm neck (i.e. the aortic segment between the lowest renal artery and the proximal extent of the aortic dilatation).



- See page 71 of the Gore Vascular/Endovascular Surgery Combat Manual in the additional resources (?@sec-additionalresources) for a high-yield EVAR planning and sizing considerations chart.
- Video Tutorial: Endovascular Aortic Repair Preoperative Sizing with Dr. Sharif Ellozy. *This is above the level of medical students*, but for those who are interested, watch it here.

The access vessels, including the bilateral iliac and femoral arteries, must be healthy enough to allow sheaths, wires and catheters to pass through them. Calcification due to atherosclerosis can make arterial access difficult, and if vessels are so diseased as to be completely occluded, wires and devices may not be able to travel past the occlusion. Additionally, the iliac arteries must be within the appropriate diameter and length dimensions to accommodate the






Figure 3: Fusiform \quad infrarenal AAA

Figure 4: Step 4.1

Figure 5: Step 4.2

EVAR Procedure




Figure 6: Step 5

Figure 7: Step 7


chosen stent graft. Lastly, certain anatomic anomalies such as a horseshoe kidney must be evaluated to ensure the appropriateness of EVAR for repair. With recent innovations in endovascular devices and intra-operative imaging, many factors that were once contraindications to EVAR have now become considerations when planning for EVAR. This is to say, as technology advances, EVAR is increasingly possible and safe for patients with difficult anatomy and co-morbidities.


EVAR Complications


Although a more recent development relative to open surgical repair (OSR), EVAR is not without risk of complications. EVAR may be complicated by injury to access vessels such as femoral occlusion or iliac avulsion (aka, "iliac-on-a-stick"), wire trauma from the relatively stiff wires that are used to deliver the stent graft, rupture of the aneurysm, and endoleak. Endoleak is the term used to describe ongoing blood flow in the aneurysm sac after stent graft placement (e.g. from lumbar arteries or the inferior mesenteric artery, inadequate proximal or distal endograft seal, a defect in the endograft, etc.). There are various types of endoleak and each has its own management approach.

Endoleaks are classified as follows:


- Type Ia: caused by an improper **proximal** endograft seal (i.e. the top of the graft is not well opposed to the abdominal aorta and blood can flow around it into the aneurysm sac).
- Type Ib: caused by an improper **distal** endograft seal (i.e. the distal iliac limb is not well opposed to the iliac artery and blood can flow retrograde into the aneurysm sac).
- Type II: caused by retrograde blood flow into the aneurysm sac from aortic side branches such as lumbar arteries or the IMA. This is the most common type and does not require intervention provided a stable aneurysm sac size is observed on follow-up. If the sac continues to expand, the feeding vessels may need to be occluded via embolization (e.g. coil embolization).
- Type III: caused by a defect in the endograft such as a fabric tear, or disconnection of the contralateral iliac limb from the main body of the graft.
- Type IV: caused by graft wall porosity. To explain graft wall porosity, imagine filling up a pillowcase with water. Even if the pillowcase has the tightest of weaves, eventually, water will leak *through* the pillowcase.
- Type V: increase in aneurysm size with no identifiable cause.








Endoleak classifications.





Tip

Please click here to open an article by Dr. Tamer W. Kassem with a simple image depicting the endoleak classifications.

Additionally, one should monitor for abdominal compartment syndrome after repair of AAA rupture, as much of the large volume resuscitation can leak from the intravascular compartment into the interstitium of the abdominal tissues causing distention. In the event of rupture, the retroperitoneal hematoma can cause a mass effect and increase of abdominal pressures. Lastly, a well-known complication of EVAR is ischemic colitis, although it is only encountered in less than 3% of endovascular aortic repairs. This occurs when the colon becomes ischemic secondary to vascular hypoperfusion caused by a variety of quoted sources including atheroembolization and systemic hypotension. Albeit rare, ischemic colitis has a significant associated mortality risk and should be addressed promptly with bowel rest, antibiotic coverage, a flexible sigmoidoscopy, and potentially further intervention, if warranted. This complication is not limited to EVAR and has been identified postoperatively with the same incidence in OSR.

Open Surgical Repair (OSR)

Open surgical repair (OSR) remains the traditional means of repairing AAA and is still the most appropriate modality when EVAR is not feasible for anatomic reasons. In the elective setting, the ideal patient is physiologically fit enough for OSR as it is a procedure with the potential for significant morbidity and a lengthy recovery.


OSR Considerations and Planning

OSR can be conducted via a transperitoneal or a retroperitoneal exposure. Other decision points for open repair include the following:

- Location of the proximal clamp: A clamp must be placed on the aorta in a location free of significant disease, with enough clearance below it to allow space for suturing the proximal anastomosis.
- Configuration of the proximal anastomosis: Anastomoses performed for an eurysmal disease are sewn in an end-to-end fashion, whereas aortas being reconstructed for occlusive disease can be performed either end-to-end or end-to-side.
- Location of the distal anastomosis: the end of the graft can be sewn to either the distal aorta, iliac arteries, or femoral arteries depending on the extent of the native vessel that needs to be excluded and the distal location where the vessel is healthy enough to clamp and receive sutures (e.g. is not heavily calcified).

- Management of the inferior mesenteric artery when an end-to-end anastomosis is created: The IMA can be ligated If collateral flow is adequate or reimplanted into the graft if collateral flow is inadequate.
- Management of aberrant intraperitoneal anatomy and pathology: Examples include adhesions or scarred tissue due to previous abdominal surgery, accessory renal arteries, horseshoe kidney, history of kidney transplant, and previous aortic replacement, among many other possibilities.

Depending on the urgency of repair, many of these considerations might only be entertained intraoperatively.



Open Aortic Repair (OAR) with suprarenal clamp.

OSR Complications

Complications related to OSR differ from those of EVAR, owing greatly to the open abdominal exposure and the lack of intraluminal wire manipulation. Just as with most other laparotomies or open retroperitoneal dissections, incisional pain can be significant and cause at lectasis or other respiratory compromise. Additionally, ileus and wound infections can impair the return of proper bowel function. Similar to EVAR, is chemic colitis can manifest after OSR, particularly in the setting of rupture.

Outcomes

Much experience has been gained since the first EVAR in 1990 and ample research has been dedicated to the comparison of EVAR to OSR. Multiple trials have demonstrated an early survival advantage for EVAR over OSR that tends to diminish over time. Certainly, the perioperative period for EVAR is notable for reduced morbidity and mortality relative to OSR with length of hospital stay significantly longer in those undergoing OSR. However, this early benefit is often balanced by a necessity for life-long follow-up including imaging at regular intervals and a tendency for re-interventions during the surveillance period (e.g. to manage endoleaks). Hence, the risk-benefit ratio must be weighed individually for each patient.

Contemporary management of abdominal aortic aneurysms is most often undertaken with EVAR. Endovascular repair has a third of the peri-operative all-cause mortality attributed to open repair (1.6% vs 4.8%, respectively). This survival benefit tends to dissipate by three to four years post-operatively. Moreover, the overall complication rate of EVAR in the perioperative period is up to 10%.

Among patients who undergo open abdominal aortic aneurysm repair, mortality in the immediate post-operative period can be as high as 5%. Of patients who survive, later death is typically related to cardiovascular disease burden. Complications arise in 9% of patients and most often include myocardial infarction, respiratory insufficiency, pneumonia, acute kidney injury, and ischemic colitis. Late Infection of the graft is possible, is challenging to manage, and occurs in less than 1% of patients.

Surveillance

The Society for Vascular Surgery guidelines suggest that for patients found to have an abdominal aortic aneurysm, surveillance imaging should occur at intervals specific to the maximum diameter of their aneurysm.

- For patients with abdominal aortic aneurysms between 3.0 and 3.9cm in diameter, CT or ultrasound should be undertaken every 3 years.
- Patients with abdominal aortic aneurysms between 4.0 and 4.9cm in diameter, imaging is indicated every 12 months.
- When the aneurysm reaches a maximum diameter of 5.0 cm, imaging with CT is recommended at intervals of every 6 months.

After the abdominal aortic aneurysm has been repaired, ongoing surveillance is warranted to monitor for post-repair complications and the need for reintervention. As mentioned, endovascular repairs can develop endoleaks over time, which may necessitate another intervention if

the aneurysm sac is expanding. Therefore, patients who have undergone EVAR should undergo CTA at one month and 12 months post-operatively. A six-month scan is indicated in the presence of a persistent endoleak observed on the one-month scan. If the endograft is in a good position without complications, yearly interval follow-up with concomitant imaging is warranted. Open repairs can similarly develop problems that require repair, albeit with less frequency, and hence need surveillance imaging every five years.

Teaching Case

Scenario

A 73-year-old male with a history of atrial fibrillation, diabetes, hypertension, ongoing tobacco use, hyperlipidemia, chronic kidney disease, and peripheral arterial disease undergoes a screening ultrasound of the abdominal aorta. The ultrasound shows a 5.6cm infrarenal fusiform aortic aneurysm extending into the proximal common iliac arteries. His primary care physician refers him to the vascular surgery clinic, where he presents without subjective complaints. CTA is recommended to assess the patient's anatomy.

Exam

HEENT: No carotid bruits.

Cardiac: Irregularly irregular, no murmurs.


Pulmonary: Clear to auscultation in all lung fields.

Abdominal: Obese, soft, non-tender, no pulsatile masses felt.

Vascular: Palpable femoral pulses, multiphasic pedal signals bilaterally.

Imaging

CTA Abdo/Pelvis



CTA Report: 5.6cm infrarenal bilobed aortic aneurysm with bilateral accessory renal arteries inferior to main renal arteries. 14mm infrarenal neck cranial to aneurysm. Widely patent iliac systems with non-diseased femoral arteries.

Discussion Points

N.B. There is no prepared answers for the questions below.

However, we feel this chapter contains all the necessary information to answer the questions. If not, please let us know!

1. What recommendations for medical management should be made to optimize this patient prior to surgical intervention?

- 2. What is the important information to ascertain from history, physical exam and imaging that can help you determine the best modality of treatment.
- 3. With what urgency should the patient be scheduled for operative repair?
- 4. What kind of repair would you suggest for this patient and why? Open or endovascular?
- 5. What complications are possible after an endovascular repair of the abdominal aorta? After an open repair?
- 6. Following repair, how should the patient be surveilled? With what frequency?
- 7. If the patient's aneurysm was 4.6cm in greatest transverse diameter, how often should he undergo imaging and follow-up?
- 8. What aortopathies and collagen vascular diseases would convince you to repair the aneurysm at a size less than 5.5cm?

Key Articles

- 1. Chaikof EL, Dalman RL, Eskandari MK, Jackson BM, Lee WA, Mansour MA, Mastracci TM, Mell M, Murad MH, Nguyen LL, Oderich GS, Patel MS, Schermerhorn ML, Starnes BW. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. J Vasc Surg. 2018 Jan;67(1):2-77.e2. doi: 10.1016/j.jvs.2017.10.044. PMID: 29268916.(Chaikof et al. 2018)
- 2. Lederle FA, Wilson SE, Johnson GR, Reinke DB, Littooy FN, Acher CW, Ballard DJ, Messina LM, Gordon IL, Chute EP, Krupski WC, Busuttil SJ, Barone GW, Sparks S, Graham LM, Rapp JH, Makaroun MS, Moneta GL, Cambria RA, Makhoul RG, Eton D, Ansel HJ, Freischlag JA, Bandyk D; Aneurysm Detection and Management Veterans Affairs Cooperative Study Group. Immediate repair compared with surveillance of small abdominal aortic aneurysms. N Engl J Med. 2002 May 9;346(19):1437-44. doi: 10.1056/NEJMoa012573. PMID: 12000813.(Lederle et al. 2002)
- 3. Lederle FA, Kyriakides TC, Stroupe KT, Freischlag JA, Padberg FT Jr, Matsumura JS, Huo Z, Johnson GR; OVER Veterans Affairs Cooperative Study Group. Open versus Endovascular Repair of Abdominal Aortic Aneurysm. N Engl J Med. 2019 May 30;380(22):2126-2135. doi: 10.1056/NEJMoa1715955. PMID: 31141634.(Lederle et al. 2019)
- 4. van Schaik TG, Yeung KK, Verhagen HJ, de Bruin JL, van Sambeek MRHM, Balm R, Zeebregts CJ, van Herwaarden JA, Blankensteijn JD; DREAM trial participants. Long-term survival and secondary procedures after open or endovascular repair of abdominal aortic aneurysms. J Vasc Surg. 2017 Nov;66(5):1379-1389. doi: 10.1016/j.jvs.2017.05.122. Erratum in: J Vasc Surg. 2018 Feb;67(2):683. PMID: 29061270.(Schaik et al. 2017)

- 5. Parodi JC, Palmaz JC, Barone HD. Transfemoral intraluminal graft implantation for abdominal aortic aneurysms. Ann Vasc Surg. 1991 Nov;5(6):491-9. doi: 10.1007/BF02015271. PMID: 1837729.(Parodi, Palmaz, and Barone 1991)
- 6. Darling, R. Master Techniques in Surgery; Vascular Surgery: Arterial. 1st ed. Lippincott Williams & Wilkins, 2016. Print.(Darling and Ozaki 2016)
- 7. Qrareya M, Zuhaili B. Management of Postoperative Complications Following Endovascular Aortic Aneurysm Repair. Surg Clin North Am. 2021 Oct;101(5):785-798. doi: 10.1016/j.suc.2021.05.020. Epub 2021 Jul 30. PMID: 34537143.(Qrareya and Zuhaili 2021)
- 8. Parkinson F, Ferguson S, Lewis P, Williams IM, Twine CP; South East Wales Vascular Network. Rupture rates of untreated large abdominal aortic aneurysms in patients unfit for elective repair. J Vasc Surg. 2015 Jun;61(6):1606-12. doi: 10.1016/j.jvs.2014.10.023. Epub 2015 Feb 7. PMID: 25661721.(Parkinson et al. 2015)
- 9. Schanzer, Andres, and Gustavo S. Oderich. Management of Abdominal Aortic Aneurysms. New England Journal of Medicine 385, no. 18 (October 28, 2021): 1690–98. https://doi.org/10.1056/NEJMcp2108504.(Schanzer and Oderich 2021)



Dr. Schanzer's recent publication in the New England Journal of Medicine (NEJM) is particularly high yield. However, as a NEJM publication, the article is behind a paywall. We hope you are able to access this paper via your institution's library.

Additional Resources

Audible Bleeding Content

- Audible Bleeding Exam Prep: AAA Chapter
- Audible Bleeding has an episode covering the IMPROVE trial. Listen to it below and find additional information here, or find the episode wherever you listen to podcasts.
- Holding Pressure Case Prep: EVAR. Listen to it here, or find the episode wherever you listen to podcasts. Be sure to take a quick look at the shownotes.
- VSITE Review AAA. Listen to it below and find additional information here, or find the episode wherever you listen to podcasts.
- Video Tutorial: Dr. Kristina Giles "Decision-Making and Treatment: Elective Infrarenal and Juxtarenal AAA." Watch it here.

•	Video Tutori	al: Open	AAA	repair:	How	I Do It	with	Dr.	Ashlee	Vinyard	and	Dr.	John
	Eidt. Watch	it here.											


•	Vascular Origin Stories:	Bridging the Gap	- The Fabric	of Aortic Repair	. Listen to it
	below and find addition	al information here	, or find the e	episode wherever	you listen to
	podcasts.				

Web:	sites
------	-------

• TeachMe Surgery: Abdominal Aortic Aneurysm

Gore Combat Manual

The Gore Medical Vascular and Endovascular Surgery Combat Manual is an informative and entertaining read intended as a vascular surgery crash course for medical students, residents, and fellows alike. Highly accessible with a thoughtfully determined level of detail, but lacking in learning activities (e.g. questions, videos, etc.), this resource is a wonderful complement to the APDVS eBook.



Operative Footage

Developed by the Debakey Institute for Cardiovascular Education & Training at Houston Methodist. *YouTube account required as video content is age-restricted.* Please create and/or log in to your YouTube account to have access to the videos.

Endovascular Aortic Repair (EVAR)

Open Surgical Repair (OSR) of a AAA, Transperitoneal Access

EVAR Explantation with Aortic Reconstruction

82

Lower Extremity Arterial Disease

Claudication

Nakia Sarad, DO, MS, Ezra Schwartz, MD, CM, MS, Marvin Chau, MD, MS, Shannon McDonnell, BS, Bernadette Aulivola, MD, FACS, Kelly Kempe, MD, RPVI, FACS, Kaled Diab, MD, and Lindsey Korepta, MD, FACS.

Note

By the end of this chapter, students will:

- Explain the definition, etiology, and pathophysiology of peripheral arterial disease and its relation to claudication.
- Identify the risk factors associated with claudication.
- Recognize the clinical manifestations and symptoms of claudication.
- Summarize the underlying vascular anatomy and physiology relevant to claudication.
- Outline the diagnostic approach and tests used to evaluate claudication.
- Summarize evidence-based outcomes of the management of claudication.

Key Facts

- 1. Peripheral arterial disease (PAD), also known as peripheral vascular disease (PVD), is a condition characterized by atherosclerosis or narrowing of the arteries that supply blood to the extremities.
- 2. PAD affects approximately 202 million individuals worldwide.
- 3. Major risk factors for PAD include smoking, diabetes, and advanced age.
- 4. Claudication is the mildest symptomatic manifestation of peripheral arterial disease.
- 5. Claudication is defined as **reproducible** discomfort in a specific muscle group, typically in the lower extremities, that is **induced by exercise and relieved by rest.**
- 6. Patients with claudication have an overall 5-year mortality risk of 10-15%.
- 7. Approximately 2-3% of claudicants (i.e. patients experiencing claudication) will develop limb-threatening ischemia.

8. Claudication is managed conservatively with lifestyle modifications including exercise regimen, dietary changes, smoking cessation, and medical optimization to control hypertension, hyperlipidemia, and or diabetes.

Claudication Slide Deck

Please find the slide deck corresponding to this eBook chapter here.

•

How We Suggest Using the Pre/Post Questions

The pre/post questions are listed below. They are all multiple choice questions with a single right answer. To best guide your learning, we have hidden the answers in a collapsible menu. Before reading the chapter, we suggest giving the questions a try, noting your answers on a notepad. After reading the chapter, return to the questions, reevaluate your answers, and then open the collapsible menu to read the correct answer and discussion. Do not fret if you have difficulty answering the questions before reading the chapter! By the end of the chapter, we are certain you will have covered the knowledge necessary to answer the questions. There will be a teaching case at the end of the chapter. This is another opportunity to exercise your new knowledge!

Pre/Post Questions

Case Based Questions

- 1. A 70-year-old male, who has a history of heavy smoking (100 pack years), presents to the clinic for claudication-like symptoms. What level of infrainguinal arterial occlusive disease is he most at risk for developing based on this history?
- A. Superficial femoral artery disease.
- B. Deep femoral artery disease.
- C. Anterior tibial artery disease.
- D. Popliteal artery disease.
- E. Posterior tibial artery disease.

Answer

A. Superficial femoral artery disease.

Discussion: The prevalence of peripheral artery disease in the general population is be-

tween 3-10%, and 15-20% in those over the age of 70. Risk factors for peripheral arterial disease (PAD) include age, male gender, hypertension, diabetes, smoking, dyslipidemia, family history and homocysteinemia. Heavy smokers have a propensity to develop superficial femoral artery disease (A). Patients with a history of diabetes have a tendency to develop infrapopliteal disease (C, E). Patients of nonwhite ethnicity and those with a history of chronic renal insufficiency are also at increased risk.

2. A 67-year-old female is being evaluated in the outpatient clinic for lower extremity pain associated with walking. Using a manual blood pressure cuff and hand-held Doppler, the following measurements are obtained:

Artery	Pressure (mm Hg)	
	Right	Left
Brachial	150	120
Dorsalis pedis (DP)	165	120
Posterior tibial (PT)	150	108

What are the ankle-brachial indices (ABIs) of her lower extremities?

- A. Right ABI 1.1; left ABI 0.8
- B. Right ABI 1.0; left ABI 1.0
- C. Right ABI 1.2; left ABI 0.9
- D. Right ABI 1.4; left ABI 1.0
- E. Right ABI 1.1; left ABI 0.7

Answer

A. Right ABI 1.1; left ABI 0.8

Discussion: Performing an ankle-brachial index (ABI) requires an appropriately sized manual blood pressure cuff and a hand-held, continuous-wave Doppler probe. The patient should be in the supine position and allowed to rest before the examination begins. The blood pressure cuff is placed above the ankle. The Doppler probe is used to locate the dorsalis pedis (DP) or posterior tibial (PT) pulse. The pressure cuff is inflated until the Doppler signal is obliterated, and then the cuff is then slowly released. The pressure at which the Doppler signal returns is the ankle pressure. The same process is repeated for both legs (DP and PT signals) and both arms. The arm pressure must be measured with the Doppler probe and the manual blood pressure cuff. To calculate an ABI, the higher ankle pressure (DP or PT) is divided by the highest arm pressure (left or right). Note that the higher arm pressure is used to calculate the ABI for both legs. For this specific

case, the highest arm pressure is 150. To calculate on the right, 165/150=1.1, on the left 120/150=0.8. A normal ABI is 0.9 to 1.2. Patients with claudication usually have an ABI ranging from 0.5 to 0.9, and those with ischemic rest pain or tissue loss usually have an ABI less than 0.5. Sabiston Textbook of Surgery: The Biological Basis of Modern Surgical Practice, 20th ed. Ch. 62: Peripheral Arterial Disease. Norgren L, Hiatt WR, Dormandy JA, Nehler MR, Harris KA, Fowkes FGR. Inter-society consensus for the management of peripheral arterial disease (TASC II). J Vasc Surg. 2007;45:S5-S67.

- 3. A 70-year-old male with poorly controlled diabetes and no smoking history presents to the clinic for an ischemic toe ulcer. What level of infrainguinal arterial occlusive disease is he most at risk for developing based on this history?
- A. Superficial femoral artery disease
- B. Deep femoral artery disease
- C. Iliac artery disease
- D. Aortic disease
- E. Tibial artery disease

Answer

E. Tibial artery disease.

Discussion: Risk factors for peripheral arterial disease include age, male gender, hypertension, diabetes, smoking, dyslipidemia, family history, and homocysteinemia. Individuals who are heavy smokers tend to develop superficial femoral artery disease, whereas those with diabetes and renal failure tend to develop infrapopliteal disease. Of the answers listed, only the tibial artery is infrapopliteal.

The following prompt will be used for questions 4, 5, and 6.

A 65-year old male with a history of hypertension, hyperlipidemia, and tobacco abuse presents to the vascular surgery clinic with complaints of pain in his legs. Patient states that the pain is crampy in quality, mostly in his calf muscles, it starts after he walks 4 blocks, and is worse on the right as compared to the left. He notices that when he stops walking, the pain resolves. This pain is keeping him from being able to participate in social events and spend quality time with his family.

- 4. What is the name of the symptom the patient is experiencing?
- A. Neuropathy
- B. Claudication
- C. Ischemic rest pain

- D. Bilateral sciatica
- E. Normal symptoms

Answer

B. Claudication

Discussion: Claudication is defined as reproducible pain or cramping of the lower extremities that occurs with ambulation and significantly improves or resolves at rest. Claudication can occur in any muscle group distal to the area of arterial stenosis and/or occlusion. As such, the calf muscles are the most common anatomic location to experience pain as they are distal in the lower extremity. As the patient does not report pain at rest, they do not have ischemic rest pain. Neuropathic pain or bilateral sciatica and claudication can be confused. Neuropathic pain is shocking in quality whereas claudication is describes as a cramp-type pain. Furthermore, claudication is absent at rest, triggered by exercise, and relieved by a return to rest. This pattern helps distinguish claudication from neuropathic pain. This patient also does not have normal pathology, as one should not have pain on ambulation and needs to be further evaluated.

- 5. What would be the first step in confirming the diagnosis for this patient?
- A. CT Angiography (CTA)
- B. Doppler probe
- C. Duplex Ultrasound
- D. Angiogram
- E. Ankle-Brachial Index and Pulse Volume Recording (ABI/PVR)

Answer

E. ABI/PVR

Discussion: Ankle Brachial Index (ABI) is a non-invasive method that measures the ratio of blood pressure in the lower legs to the blood pressure in the arms. Abnormal ABIs have been shown to have high specificity with identifying PAD. Patients who have claudication usually have ABI ranges from 0.5 to 0.9. Pulse volume recordings (PVR) use pressure cuffs to record the shape of a pulse. The type of waveform provides important information about the characteristics of blood flow. Waveforms can be used to diagnose and monitor arterial diseases, assess the severity of stenosis, and determine the effectiveness of treatment interventions. This patient likely has claudication based on the history and symptoms. The first non-invasive testing to confirm the diagnosis would be ABI/PVR. While a CTA would provide helful information with regards to arterial anatomy and the extend of calcification, it cannot provide the physiologic information that one would ob-

tains an ABI/PVR. Furthermore, CTA requires contrast and radiation exposure. While used in measuring an ABI, a Doppler probe is not nearly as sensitive in identifying PAD and does not provide data helpful in creating a management plan. A Duplex Ultrasound would provide physiologic information as to the burden of disease. However, it is highly operator dependent and requires more resources. It may be considered as a second line investigation. While an angiogram is the gold-standard, it is an invasive technique that requires the use of contrast. As such, it is not a first line investigation.

- 6. What would be the first step in treating this patient's symptoms?
- A. Angiogram to identify targets for open surgical bypass.
- B. Angiogram with angioplasty/stent of any occluded vessels.
- C. Prescribe Cilostazol.
- D. Ensure medication optimization for his hypertension and hyperlipidemia, smoking cessation counseling, and exercise regimen.
- E. No treatments at this time, continue to monitor and come back to clinic in 3 months.

Answer

D. Ensure medication optimization for his hypertension and hyperlipidemia, smoking cessation counseling, and exercise regimen.

Discussion: The mainstay of treatment for claudication is conservative management, including medical optimization and lifestyle modification. The patient has hypertension and hyperlipidemia. He should be on appropriate antihypertensive to control the blood pressure. With regard to medical optimization, it is recommended that patients start on antiplatelet therapy (Aspirin 81 mg daily) and a high dose statin. Lifestyle modifications include smoking cessation, exercise training to develop collateral vessels, and diet modification to lower cholesterol intake. An appropriate exercise regimen is defined by at least 30 minutes of supervised ambulation/exercise at least 3 times per week. Patients should be followed closely with yearly ABIs by both Primary Care Physicians (PCPs) and Vascular Surgeons for progression of disease. While an angiogram allows for simultaneous diagnosis and treatment of PAD, it is an invasive technique that requires the use of contrast. Angiogram may be considered if conservative management fails. Cilostazol is an antiplatelet that has been shown to have beneficial effects for patients with intermittent claudication, however, it is not a first-line medication. Lastly, while follow-up is important in the management of PAD, conservative management should be initiated as soon as the diagnosis of PAD is made.

Introduction



Tip

The term peripheral arterial disease (PAD) is often used interchangeably with peripheral vascular disease (PVD) when referring to arterial blockages.

Claudication is defined as cramping, pain, and/or discomfort in the hip, thigh, or calves that occurs with ambulation/exercise, improves shortly after cessation of activity, and recurs after a predictable amount of ambulation/exercise.

Claudication is a symptom of peripheral arterial disease (PAD). Therefore, it is important to first understand PAD before addressing claudication. PAD is a chronic condition of arterial insufficiency due to the build-up of atherosclerotic plaque (or fatty deposits) in the arteries. PAD affects 8.5 million adults in the U.S. and 202 million adults worldwide, with approximately 15% of the population aged greater than 70 years old suffering from PAD. Risk factors for PAD include increasing age, hypertension, hyperlipidemia, renal disease, diabetes, and smoking. The risk factors for claudication are the same risk factors described for PAD, including: advanced age, race (non-hispanic blacks), male gender, smoking, diabetes mellitus, dyslipidemia, hypertension, renal insufficiency, and elevated homocysteine levels (hypercoagulable state). PAD has various distributions in the body including one or a combination of aortoiliac, femoral, and infrapopliteal disease. PAD severity falls along a spectrum. Some patients may be completely asymptomatic while others have nonhealing, chronically infected wounds requiring later limb amputation. Vascular claudication is the mildest symptomatic manifestation of the PAD spectrum.

Etiology

Prior to understanding the etiology of claudication and peripheral arterial disease. It is important to review the anatomy of the arterial blood flow of the lower extremity.

Anatomy

The arterial anatomy of the lower extremity is a complex network of arteries that supply blood to the legs and feet. Please find an <u>image here</u> and a description below of the major arteries from the common femoral artery down to the tibioperoneal trunk, anterior tibial, and dorsalis pedis arteries:

- Common Femoral Artery (CFA): The common femoral artery is a large, thick-walled artery located in the groin. It is a continuation of the external iliac artery and bifurcates into the deep femoral artery (profunda femoris) and the superficial femoral artery.
- Superficial Femoral Artery (SFA): The superficial femoral artery is the largest branch of the common femoral artery. It descends along the anterior and medial aspect of the thigh and enters the adductor canal. As it descends, it eventually becomes the populated artery at the level of the knee joint.
- Profunda Femoris (Deep Femoral) Artery (PFA): The profunda femoris artery, also known as the deep femoral artery, is a significant branch of the common femoral artery. It typically arises about 1 to 2 cm below the inguinal ligament. The profunda femoris artery primarily supplies the muscles of the posterior and deep thigh, including the quadriceps muscles. It is also an important collateral pathway for blood flow in cases of arterial obstruction or occlusion in the lower extremity distal to the SFA/PFA bifurcation.
- **Popliteal Artery:** The popliteal artery is located behind the knee joint and is a continuation of the superficial femoral artery. It provides blood supply to the muscles and structures in the popliteal fossa and gives rise to several branches, including the anterior tibial, posterior tibial, and peroneal arteries.
- Anterior Tibial Artery (AT): The anterior tibial artery is the first branch off of the popliteal artery. It is NOT part of the tibioperoneal trunk (only includes the PT and peroneal arteries. The AT passes from behind the knee through the interosseous membrane of the leg and descends through the anterior compartment of the leg. It can be visualized as the most lateral tibial vessel during lower extremity angiogram run-offs. The AT runs along the front of the tibia and crosses the ankle joint to become the dorsalis pedis artery, supplying blood to the dorsum (top) of the foot.
- Posterior Tibial Artery (PT): The posterior tibial artery is another branch of the popliteal artery. Once the AT tibial branches from the popliteal, the remaining vessel is termed the tibeoperoneal trunk. This trunk splits to give off the posterior tibial artery (PT) and the peroneal artery. The PT descends posteriorly in the leg and passes behind the medial malleolus (the bony prominence on the inner side of the ankle). It can be palpated on the medial aspect of the foot behind the medial malleolus. It continues into the sole of the foot and forms the plantar arch, contributing to the arterial supply of the sole of the foot and toes.
- **Peroneal Artery:** The peroneal artery, also known as the fibular artery, is the third major branch of the popliteal artery. As mentioned, it originated from tibioperoneal trunk along with the PT. The peroneal artery runs along the lateral side of the leg and supplies the lateral compartment muscles. In the ankle region, it anastomoses with the branches of the posterior tibial artery, providing collateral circulation.
- Dorsalis Pedis Artery (DP): The dorsalis pedis artery is a continuation of the anterior tibial artery. It extends along the anterior aspect of the ankle and runs atop the dorsum (top) of the foot. It can be palpated on the dorsum of the foot just lateral to the tender of extensor hallucis longus. It plays a crucial role in supplying blood to the dorsal surface of the foot and the toes. The presence and palpability of the dorsalis pedis pulse are

often evaluated as part of a clinical assessment to determine arterial perfusion in the foot.

In summary, the common femoral artery is the arterial origin of the leg. It gives rise to the superficial femoral artery, and subsequently, the popliteal artery. The popliteal artery then divides into the anterior tibial, posterior tibial, and peroneal arteries, each supplying specific regions of the leg and foot. Understanding this anatomy is crucial for diagnosing and treating arterial diseases and vascular disorders in the lower extremities.



Pip

For a short, comprehensive, although advanced video lecture reviewing normal and diseased lower limb arterial anatomy, please see the video below.

Note

Here is a review of the lower extremity vascular anatomy and here are two sample diagnostic angiograms of these vessels (Sample 1, Sample 2).

Process of Atherosclerosis

As previously mentioned, atherosclerosis is a fundamental process that underlies the development of peripheral vascular disease in the lower extremities. The pathological process is precisely the same as with carotid disease leading to increasing stroke risk and coronary artery disease leading to increased risk of cardiac events.

Atherosclerosis is a progressive, inflammatory condition characterized by the accumulation of fatty deposits, cholesterol, inflammatory cells, and fibrous tissue within the inner layers of arteries, forming plaques. The process typically begins with damage or dysfunction of the endothelial lining of the arterial walls. This can be caused by risk factors such as hypertension, smoking, or high cholesterol levels. In response to endothelial injury, low-density lipoproteins (LDL) or "bad" cholesterol molecules penetrate the arterial wall and become oxidized. These oxidized lipids trigger an inflammatory response. The body's immune system responds to the inflammatory signals by sending white blood cells (macrophages) to the site. These cells engulf the oxidized lipids and form foam cells within the arterial wall. The inflammation leads to the proliferation of smooth muscle cells. Over time, the combination of lipids, immune cells, and smooth muscle cell proliferation results in the formation of atherosclerotic plaques. These plaques narrow the arterial lumen, restricting blood flow. The build-up of plaques can lead to stiffening and loss of elasticity in the arterial walls, further diminishing the artery's ability to dilate and maintain adequate blood flow.

As the arterial narrowing progresses, blood flow to the downstream tissues (e.g., muscles, nerves, and skin in the lower extremities) becomes compromised. This can lead to ischemia, which manifests as claudication – a symptom characterized by intermittent pain, cramping, and fatigue in the muscles during physical activity that resolves once at rest. In severe cases, atherosclerosis can lead to critical limb threatening ischemia (CLTI), a condition marked by severe arterial blockages and tissue damage. CLTI can result in non-healing ulcers, gangrene, and, in the absence of intervention, limb amputation. Atherosclerotic plaques are also prone to rupture or ulceration, exposing their contents to the bloodstream. This can lead to the formation of blood clots (thrombosis) or embolization of plaque fragments to smaller arteries, causing acute blockages that may lead to acute limb ischemia (ALI). CLTI and ALI will be discussed in more detail in the subsequent PAD chapters.

Note

The following lipid levels are considered to be "good" in healthy people:

- Total cholesterol: Levels below 200 mg/dL (5.2 mmol/L)
- LDL cholesterol: Levels below 130 mg/dL (3.4 mmol/L)
- HDL cholesterol: Levels above 40 mg/dL (1 mmol/L) in men and above 50 mg/dL (1.3 mmol/L) in women

The American Diabetes Association (ADA) standards of care for diabetes recommend that statin therapy should be initiated in individuals with diabetes and other atherosclerotic cardiovascular disease (ASCVD) risk factors.

Please see the ASCVD risk calculator algorithm:

- For patients ages 40 years and older with diabetes: Use moderate-dose statin therapy.
- For patients ages 40 to 75 with diabetes and a higher cardiovascular risk (including those with one or more ASCVD risk factors): Use high-intensity statin therapy to reduce LDL cholesterol by 50% of baseline and target an LDL cholesterol goal of <70 mg/dL.
- For patients with diabetes who have already had an ASCVD event: High-intensity statin therapy is recommended to target an LDL cholesterol reduction of 50% from baseline and an LDL cholesterol goal of <55 mg/dL.

PAD Risk Factors

All these risk factors increase rates of atherosclerosis and start to limit the blood flow to the lower extremities, attributing to claudication.

• Smoking: Smoking is a well-established major risk factor for PAD. It contributes to PAD through multiple mechanisms, including endothelial dysfunction, oxidative stress, inflammation, and platelet activation. The nicotine in tobacco constricts blood vessels

- and promotes the formation of atherosclerotic plaques. Moreover, smoking cessation is the most effective intervention in reducing the progression of PAD.
- Diabetes Mellitus: Diabetes is closely associated with PAD due to hyperglycemiainduced endothelial dysfunction and microvascular damage. It accelerates atherosclerosis by promoting lipid abnormalities, systemic inflammation, and the development of neuropathy, which can mask or delay the recognition of PAD symptoms.
- **Hypertension:** Hypertension is a major contributor to PAD. Elevated blood pressure damages arterial walls, rendering them more susceptible to atherosclerosis. Hypertension also increases the risk of thrombosis and clot formation, further compromising arterial perfusion.
- **Hyperlipidemia:** High levels of low-density lipoprotein (LDL) cholesterol are a known risk factor for PAD. Elevated LDL cholesterol leads to the accumulation of lipid deposits in arterial walls, initiating the atherosclerotic process. Dyslipidemia management, particularly the reduction of LDL cholesterol, is essential in PAD prevention and management.
- Age: Age is an independent risk factor for PAD. As individuals age, atherosclerotic processes accumulate, contributing to a higher likelihood of PAD development.
- Family History: Genetic predisposition can influence an individual's susceptibility to PAD. A family history of vascular disease, particularly atherosclerosis, can confer a higher risk.
- Obesity: Obesity is one of the multifactorial contributors to the formation of PAD. Excess adipose tissue is associated with insulin resistance, chronic inflammation, and altered lipid profiles. These factors contribute to atherosclerosis and PAD development.
- Physical Innactivity: A sedentary lifestyle not only contributes to obesity but also reduces overall vascular health. Regular physical activity, conversely, is associated with better arterial function and can mitigate PAD risk.
- **Diet:** A diet rich in saturated and trans fats, along with a low intake of fruits and vegetables, promotes the development of atherosclerosis. In contrast, diets emphasizing heart-healthy components, such as whole grains, fruits, vegetables, and lean protein, are associated with reduced PAD risk.
- Metabolic Syndrome: Metabolic syndrome, which encompasses a cluster of conditions such as obesity, hypertension, hyperglycemia, and dyslipidemia, is a potent risk factor for PAD. The interplay of these factors significantly increases the risk of atherosclerosis and impaired arterial perfusion.
- Gender and Menopause: Men generally exhibit a higher prevalence of PAD, although the risk in women increases post-menopause. Reduced estrogen levels post-menopause contribute to endothelial dysfunction and alterations in lipid profiles, increasing susceptibility to PAD.

- Race and Ethnicity: Certain racial and ethnic groups, such as African Americans, are more prone to developing PAD. Genetic and environmental factors may contribute to these disparities.
- Chronic Kidney Disease (CKD): Reduced kidney function is associated with PAD, partly due to endothelial dysfunction and systemic inflammation. CKD patients are at increased risk of atherosclerosis, uncontrolled hypertension, thrombotic complications, and progressive PAD.

Vessel Collaterization

Patients with peripheral arterial disease (PAD) often develop collateralization of blood vessels as a compensatory mechanism to improve perfusion to tissues affected by reduced blood flow. These new blood vessels, or collaterals, bypass or supplement the obstructed arteries and combat oxygen starvation of the tissues through "collateral circulation." The main drivers for the development of collateral vessels in PAD are as follows:

- Ischemia and Hypoxia: In PAD, atherosclerosis narrows and sometimes occludes the arteries that supply blood to the limbs. The resulting reduced blood flow causes tissue ischemia (insufficient blood supply) and hypoxia (insufficient oxygen supply) to the tissues distal to the narrowing or occlusion. To combat this, the body initiates a response to create alternate pathways for blood flow.
- Hypoxia-Inducible Factors (HIFs): When cells are deprived of oxygen (as in the case of PAD), they activate hypoxia-inducible factors (HIFs). HIFs are transcription factors such as vascular endothelial growth factor (VEGF) and fibroblast growth factor (FGF). These factors promote the expression of genes involved in angiogenesis (the formation of new blood vessels) and arteriogenesis (the remodeling and enlargement of existing collateral vessels) through the stimulation of endothelial cells. Endothelial cells line blood vessels and play a central role in angiogenesis. They respond to these growth factors enabling endothelial cell proliferation and migration, which are essential for the formation of new capillaries in the direction of the hypoxic tissue.
- Vasodilation: When blood vessels are exposed to reduced oxygen levels, they often dilate in an attempt to increase blood flow. This vasodilation can help mobilize and enhance the development of collateral vessels.
- Angiogenesis and Arteriogenesis: Angiogenesis involves the formation of small capillaries, while arteriogenesis involves the enlargement and maturation of existing collateral vessels into larger arteries. These processes are driven by growth factors and signaling molecules like vascular endothelial growth factor (VEGF), fibroblast growth factor (FGF), and various cytokines. As stated previously, they promote the proliferation and migration of endothelial cells and smooth muscle cells, leading to the formation of new blood vessels and the remodeling of existing ones.

- Shear Stress: As blood flows through collateral vessels, it exerts shear stress on the vessel walls. Shear stress is a mechanical force that can trigger the release of molecules like nitric oxide, which further stimulates vessel growth and remodeling.
- Physical Activity: Exercise and physical activity can also stimulate collateralization by increasing shear stress on the vessel walls, enhancing collateral vessel growth. Furthermore, physical activity increases oxygen demand of the active tissues creating a relative hypoxic environment, leading to collateralization/angiogenesis/arteriogenesis as described above. This is why dedicated exercise is an important aspect of treatment for claudication.

The development of collateral circulation in PAD is a gradual process that occurs over time in response to the chronic reduction in blood flow. While collateralization can help improve perfusion to some extent, it may not fully compensate for the lost function of the original, diseased arteries. In advanced stages of PAD, when the arterial obstruction is severe, collateralization may become insufficient, leading to continued symptoms of intermittent claudication, pain, and impaired wound healing.

Anatomic Distribution of PAD

The distribution of disease in PAD varies among different risk groups such as smokers and diabetics. Smokers tend to develop PAD predominantly in the infrainguinal arteries (CFA, SFA, PFA, popliteal artery), while diabetics tend to have higher infrapopliteal disease patterns (AT, TP, peroneal, DP). (Conte et al. 2019a)

Smoking is a major risk factor for atherosclerosis, which is the primary underlying cause of PAD. Atherosclerosis tends to affect larger and proximal arteries, including the femoral and popliteal arteries, in smokers. The buildup of atherosclerotic plaques narrows these arteries, reducing blood flow to the lower extremities and causing symptoms like claudication.

Diabetes is associated with a specific type of arterial disease known as diabetic vasculopathy or microangiopathy. This condition leads to changes in the smaller blood vessels, including the tibial arteries and peroneal artery. The tibial arteries play a critical role in supplying blood to the feet, making them susceptible to diabetic vascular complications. Therefore, the microvascular changes associated with diabetes result in a distal distribution of arterial disease (i.e. toes and feet first).

96

Diagnostics and Imaging

Measurement of the ankle-brachial index (ABI) is the primary method for establishing the diagnosis of PAD, however, a proper physical exam helps confirm the diagnosis of claudication.

Peripheral Pulse Exam

In terms of physical exam, claudicants may exhibit signs of muscle wasting with thin, dry, and hairless skin, or even ulceration. These are all signs that blood flow is reduced to the point that there are not enough nutrients to feed these tissues. If ulceration or non-healing wounds are present, the patient likely has advanced peripheral arterial disease (i.e. critical limb ischemia). In addition, a lower extremity motor/sensory exam and peripheral pulse assessment is key.

Note

Pulse exams study the presence, strength, and character of arterial blood flow and are graded on a scale from 0 to 4+. 0: non-palpable pulse 1+: diminished pulse 2+: normal pulse 3+: prominent pulse or aneurysmal. 4+: bounding pulse or aneurysmal.

Lower extremity peripheral pulse exams evaluate the dorsalis pedis (DP) and posterior tibial (PT) pulses using manual palpation. Patients with claudication may have pulse exams ranging from non-palpable to normal pulse. If patients have non-palpable pulses, then a continuous-wave Doppler probe can be used to assess blood flow.

Ankle Brachial Index (ABI)

Ankle Brachial Index (ABI) is a non-invasive method that measures the ratio of blood pressure in the lower legs to the blood pressure in the arms. Performing an ABI requires an appropriately sized manual blood pressure cuff and a hand-held, continuous-wave Doppler probe. The patient should be in the supine position and allowed to rest before the examination begins. The blood pressure cuff is placed above the ankle. The Doppler probe is used to locate the dorsalis pedis (DP) or posterior tibial (PT) pulse, the cuff is inflated until the Doppler signal is obliterated, and then the cuff is then slowly released; the pressure at which the Doppler signal returns is the ankle pressure. The same process is repeated for both legs (DP and PT signals) and both arms. The arm pressure must be conducted with the manual blood pressure cuff that is placed above the target vessel and the Doppler probe that is placed below the cuff to evaluate blood flow. To calculate the ABI, the higher ankle pressure (DP or PT) is divided by the highest arm pressure (left or right).

ABI result of <0.9 has shown to have high sensitivity and specificity with identifying PAD. Patients with claudication usually have an ABI ranging from 0.5 to 0.9, and those with ischemic rest pain or tissue loss usually have an ABI less than 0.5.

Note

Below are the parameters used to diagnose PAD using ABI values.

- 1. ABI of >1.2: Calcification of arteries, non-compressible. Seen in diabetic patients.
- 2. ABI of 0.9 1.2: Normal
- 3. ABI of <0.9: Suggests arterial stenosis. (Khan, Farooqui, and Niazi 2008)
- ABI 0.7-0.9 = mild PAD
- ABI 0.4-0.7 = moderate PAD
- ABI < 0.4 = severe PAD

ABI 0.5 - 0.9: often causes claudication

ABI <0.5: often causes ischemic rest pain/tissue loss

Below, we will review how to calculate the ABI for the right and left lower extremity using the following values:

Artery	Pressure (mm Hg)	
	Right	Left
Brachial	150	130
Dorsalis pedis (DP)	145	120
Posterior tibial (PT)	150	110

For both right and left lower extremities, the highest brachial pressure is 150 and will be used for comparison for the ankle/brachial index.

On the right lower extremity, using the DP systolic pressure the ABI is 145/150 = 0.97 and using the PT systolic pressure the ABI = 150/150 = 1.0. So the Right ABI is 1.0, because we choose the highest of the two ankle pressures for the final value of the ABI.

On the left lower extremity, using the DP systolic pressure the ABI is 120/150 = 0.8 and using the PT systolic pressure the ABI = 110/150 = 0.73. So the Left ABI is 0.8, because we choose the highest of the two ankle pressures for the final value of the ABI.

For this above example, R ABI = 1.0 and L ABI = 0.8. So this patient may have some peripheral vascular disease present in his left lower extremity as his ABI is less than 0.9.



Here is a video demonstrating how to obtain and calculate an ABI.

Pulse volume recordings (PVR)

Pulse volume recordings (PVR) use pressure cuffs to record the shape of a pulse. The type of waveform provides important information about the characteristics of blood flow. These waveforms can be categorized as triphasic, bisphasic, or monophasic. Clinicians use these waveform patterns to diagnose and monitor arterial diseases, assess the severity of stenosis, and determine the effectiveness of treatment interventions.

Pulse volume recordings (PVR) use pressure cuffs to record the shape of a pulse. The type of waveform provides important information about the characteristics of blood flow. These waveforms can be categorized as triphasic, bisphasic, or monophasic. Clinicians use these waveform patterns to diagnose and monitor arterial diseases, assess the severity of stenosis, and determine the effectiveness of treatment interventions.

• Triphasic Waveform:

- A triphasic waveform consists of three phases: forward flow during systole, a brief reverse flow during early diastole, and a second forward flow component during late diastole.
- Clinical Significance: Triphasic waveforms are typically found in healthy, unobstructed arteries. They represent normal arterial blood flow. These waveforms are seen in arteries serving well-perfused tissues where resistance is low, and blood flow is unimpeded. Triphasic waveforms are commonly observed in the larger arteries of the extremities, such as the femoral and brachial arteries.

• Biphasic Waveform:

- A biphasic waveform displays two phases: an initial forward flow during systole (when the heart contracts and ejects blood) followed by brief reverse or "reflected" flow during early diastole.
- Clinical Significance: Biphasic waveforms are often associated with moderate arterial stenosis. The reverse flow component indicates some degree of obstruction. They can be observed in patients with early-stage PAD or moderate occlusions. Biphasic waveforms suggest that some blood is able to flow past the obstruction during systole but encounters resistance during diastole.

• Monophasic Waveform:

- A monophasic waveform consists of a single, continuous flow signal that moves in one direction. It typically lacks a distinct reverse or "reflected" flow signal.
- Clinical Significance: Monophasic waveforms are often seen in arteries downstream from significant stenosis or occlusion. In these cases, there is limited or no backflow during diastole (the relaxation phase of the cardiac cycle). They are commonly found in patients with advanced peripheral arterial disease (PAD) or severe arterial blockages, suggestive of reduced perfusion to the tissues served by the artery.

• Flat-line:

• A flat waveform (i.e. no wave) does not necessarily indicate no blood flow. A flat wave form simply reflects the lack of pulsatile blood flow.

Doppler

Doppler devices (handheld or otherwise) provide similar information to PVR studies. Instead of providing a visual representation of pulsatile flow, Doppler signals (sounds) are an auditory representation of flow. A triphasic pulse generates a triphasic signal (repeated sound), a biphasic pulse a biphasic signal, and a monophasic pulse a monophasic signal.



Here is a video that demonstrates how to use a doppler probe and the signals associated with each waveform type. You can learn more about doppler probes and waveforms here.

Duplex Ultrasonography

Duplex ultrasonography (US) is another non-invasive imaging modality that can evaluate a patient's blood flow by measuring the speed at which blood travels through the vessel. In this way, a patient's hemodynamics can be assessed by identifying the location and degree of stenosis. Other imaging modalities such as a CT angiogram or invasive angiograms can also evaluate blood flow to the lower extremities.

A hematologic evaluation (including CBC, fasting blood glucose, creatinine, fasting lipid profile and urinalysis) is recommended in order to identify any associated comorbidities such as diabetes, hyperlipidemia, and chronic kidney disease that may also need to be addressed.

If patients have normal ABIs and pulse exams, it is important to consider other differentials of bilateral lower extremity pain include hematologic or neurogenic claudication (pain caused by lumbosacral nerve root impingement). Neurogenic claudication is positional, often worse when walking downhill and and relieved by leaning forward (i.e. leaning on a walker).

Treatment

The mainstay of treatment for claudication is conservative management, including medical optimization and lifestyle modification. With regard to medical optimization, it is recommended

that patients start on antiplatelet therapy (Aspirin 81 mg daily) and a high dose statin. Patients without a prior history of congestive heart failure can also start on cilostozal (Pletal) to help with pain management. Patients with prior history of hypertension and diabetes should also be on appropriate antihypertensives to control their blood pressure and diabetes medications to control their blood sugar levels. Lifestyle modifications include smoking cessation, exercise training to develop collateral vessels, and diet modification to lower cholesterol intake. An appropriate exercise regimen is defined by at least 30 minutes of supervised ambulation/exercise at least 3 times per week. Patients should be followed closely with yearly ABIs by both Primary Care Physicians (PCPs) and Vascular Surgeons for progression of disease.

Below is an outline of appropriate treatment options for claudication:

Supervised Exercise Regimen

Goal of at least 30 minutes of supervised walking exercise (at least 3 times a week)

Optimization of cardiovascular risk factors:

Hypertension - Use of antihypertensive medications - Identify any underlying etiology of hypertension (ie renal artery stenosis, stressors)

Hyperlipidemia - Control of cholesterol levels - Diet/exercise modification - Use of medications: i.e. statins, anti-platelet agents

Diabetes mellitus - Control of blood glucose levels - Diet/exercise modification - Medications (i.e. antihyperglycemics)

Smoking Cessation

Pain Management

Anti-inflammatories (ie acetaminophen, ibuprofen)

Cilostazol

Outcomes

When treated appropriately, most patients with claudication will not experience limb loss. Only 1-3% of claudicants develop critical limb ischemia and the complications associated with more severe PAD. While the majority of patients (70-80%) have stable claudication symptoms, approximately 20-30% of patients experience a worsening of their claudication. In terms of overall morbidity and mortality, patients with claudication have worse overall outcomes than the general population. 20% of patients with claudication will experience either a myocardial infarction (MI) or cerebrovascular accident (CVA) within their lifetime. Moreover, patients with diagnosis of claudication have a 5-year mortality rate of 10-15% due to the comorbidities associated with vascular disease. (Conte et al. 2015a)

Teaching Case

Scenario

73 year old male with past medical history of hypertension, hyperlipidemia, and diabetes presents to clinic for pain in his lower extremities. Of note, he has smoked roughly one pack of cigarettes a day since he was 20 years old. He states that it is difficult for him to walk 3 blocks without having to stop which he finds particularly annoying. When asked where it hurts, he starts grabbing his calves. He describes an achy, cramping, and pain in his calves whenever he walks, mostly on the right side. He denies any new sores or wounds that are not healing.

Exam

HEENT: pupils equal round reactive to light, no lymphadenopathy

Cardiac: Regular rate and rhythm (RRR)

Pulmonary: Clear to auscultation in all lung fields. No accessory muscle use.

Abdominal: Soft, non-tender, non-distended, no pulsatile masses felt.

Vascular/Extremities: Warm, no edema present. Minimal hair appreciated bilaterally. Non-palpable distal pulses. Bilateral femoral pulses appreciated. No wounds or sores noted.

Imaging

ABI: Right: 0.7 Left: 0.65

Discussion Points

N.B. There is no prepared answers for the questions below.

However, we feel this chapter contains all the necessary information to answer the questions. If not, please let us know!

- 1. Please explain the pathophysiology for the patient's lower extremity pain and discomfort.
- 2. Please list the patient's risk factors for peripheral arterial disease. What is the best medical management to optimize the these risk factors?
- 3. What initial testing would you want to obtain and why?
- 4. What would be the first line of treatment for patients with symptomatic claudication after medical management?

Key Articles

- 1. Conte MS, Pomposelli FB, Clair DG, et al. Society for Vascular Surgery practice guidelines for atherosclerotic occlusive disease of the lower extremities: Management of asymptomatic disease and claudication. Journal of Vascular Surgery. 2015;61(3):2S-41S.e1. doi:10.1016/j.jvs.2014.12.009(Conte et al. 2015b)
- 2. Conte MS, Bradbury AW, Kolh P, et al. Global vascular guidelines on the management of chronic limb-threatening ischemia. Journal of Vascular Surgery. 2019;69(6):3S-125S.e40. doi:10.1016/j.jvs.2019.02.016(Conte et al. 2019b)
- 3. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the Management of Patients With Lower Extremity Peripheral Artery Disease: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. Circulation. 2017;135(12). doi:10.1161/CIR.000000000000000470(Gerhard-Herman et al. 2017a)
- Khan TH, Farooqui FA, Niazi K. Critical review of the ankle brachial index. Curr Cardiol Rev. 2008 May;4(2):101-6. doi: 10.2174/157340308784245810. PMID: 19936284; PMCID: PMC2779349.(Khan, Farooqui, and Niazi 2008)
- 5. McDermott MM, Criqui MH. Ankle-Brachial Index Screening and Improving Peripheral Artery Disease Detection and Outcomes. JAMA. 2018;320(2):143. doi:10.1001/jama.2018.8953(McDermott and Criqui 2018)

Additional Resources

Audible Bleeding Content

- Audible Bleeding Exam Prep: Medical Management and Claudication Chapter
- Caitlin W. Hicks, MD, MS Overuse of early peripheral vascular interventions in claudication. Listen to it below and find additional information here, or find the episode wherever you listen to podcasts.
- VSITE Review Peripheral Arterial Occlusive Disease (Part One). Listen to it below and find additional information here, or find the episode wherever you listen to podcasts.

Gore Combat Manual

The Gore Medical Vascular and Endovascular Surgery Combat Manual is an informative and entertaining read intended as a vascular surgery crash course for medical students, residents, and fellows alike. Highly accessible with a thoughtfully determined level of detail, but lacking in learning activities (e.g. questions, videos, etc.), this resource is a wonderful complement to the APDVS eBook.



Please see pages 85-90.

Chronic Limb Threatening Ischemia (CLTI)

Nakia Sarad, DO, MS, Ezra Schwartz, MD, CM, MS, Chaitanya Dandu, MD, Marvin Chau, MD, Shannon McDonnell, BS, Bernadette Aulivola, MD, FACS, Kelly Kempe, MD, RPVI, FACS, Kaled Diab, MD, and Lindsey Korepta, MD, FACS.

Note

By the end of this chapter, students will:

- Define chronic limb-threatening ischemia (CLTI) and summarize its etiology and pathophysiology.
- Identify the clinical features and diagnostic criteria used to diagnose CLTI.
- Elaborate on the complications and potential consequences of untreated CLTI.
- Outline a comprehensive history and physical examination to evaluate patients with suspected CLTI.
- Interpret diagnostic tests such as ankle-brachial index (ABI), Doppler ultrasound, and angiography to confirm the diagnosis.
- Develop and implement a management plan for patients with CLTI, including medical, endovascular, or surgical interventions.

Key Facts

- 1. Chronic limb-threatening ischemia (CLTI) refers to a severe and advanced form of peripheral arterial disease (PAD) characterized by chronic inadequate blood supply to the lower extremities for **greater than 2 weeks.**
- 2. CLTI presents as ischemic rest pain, tissue loss (e.g. ulcers, gangrene), or both.
- 3. CLTI is associated with amputation, impaired quality of life, and mortality.
- 4. Rutherford Classification and Fontaine Grading are classification systems developed to categorize the degree of severity of CLTI, which also aids in management.
- 5. Newer classification systems including WiFI (Wound, Ischemia, and Foot Infection)

and GLASS (Global Limb Anatomic Staging System) aim to aid in prognostication and management of CLTI.

Claudication Slide Deck

Please find the slide deck corresponding to this eBook chapter here.

How We Suggest Using the Pre/Post Questions

The pre/post questions are listed below. They are all multiple choice questions with a single right answer. To best guide your learning, we have hidden the answers in a collapsible menu. Before reading the chapter, we suggest giving the questions a try, noting your answers on a notepad. After reading the chapter, return to the questions, reevaluate your answers, and then open the collapsible menu to read the correct answer and discussion. Do not fret if you have difficulty answering the questions before reading the chapter! By the end of the chapter, we are certain you will have covered the knowledge necessary to answer the questions. There will be a teaching case at the end of the chapter. This is another opportunity to exercise your new knowledge!

Pre/Post Questions

Case Based Questions

- 1. A 45-year-old male with a history of type 2 diabetes mellitus presents to the emergency department with wet gangrene of his first, fourth, and fifth digits of his right lower extremity. He has a right dorsalis pedis and posterior tibial artery pulse. What level of amputation is appropriate in this situation?
- A. Transmetatarsal amputation
- B. Below-knee amputation
- C. Through-knee amputation
- D. Above-knee amputation

Answer

A. Transmetatarsal amputation (TMA).

Discussion: This patient requires removal of the affected toes. However, if only the toes are removed, the remaining toes are subject to ulcer formation and or infection. If 3 or more toes are removed, the stability of the foot can be compromised and thus the correct amputation for him is a transmetatarsal amputation (TMA). He has adequate distal perfusion with palpable pedal pulses, and therefore, he will likely be able to heal from the amputation. A TMA will remove infected tissue and will provide him with the greatest amputation stump length. A below-knee amputation (BKA), through-knee amputation (TKA), and above-knee amputation (AKA) are too extreme for the burden of foot infection. Furthermore, the other alternatives are associated with greater morbidity and amount of energy required to ambulate, with the degree of severity increasing the more proximal the amputation (AKA is worse than BKA which is worse that TMA). Lower extremity amputation: techniques and results. In: Cronenwett JL, Johnson KW, eds. Rutherford's Vascular Surgery. 8th ed. Philadelphia, PA: Elsevier; 2014.

- 2. A 72-year-old male with diabetes presents to his PCP's office for a nonhealing ulcer to the plantar aspect of his right foot. Laboratory results and x-ray imaging are unremarkable. On examination, you note that his pulses are nonpalpable. Doppler examination reveals monophasic signals. He has necrotic, nonviable tissue to the base of the wound. There is no erythema or evidence of abscess. The patient has not been taking antibiotics. What is the most likely cause of the non-healing in this scenario?
- A. Pressure
- B. Infection
- C. Inadequate wound care
- D. Peripheral vascular (arterial) disease

Answer

D. Peripheral vascular (arterial) disease

Discussion: This patient has nonpalpable pulses and monophasic doppler signals. Along with necrotic tissue to the wound base, these findings are consistent with peripheral artery disease (PAD). You may choose to order ankle-brachial index/pulse volume recording studies. He should be referred to a vascular specialist for further workup before any intervention is made. He does not have erythema or evidence of abscess on examination, and therefore it is not likely that primary infection is the cause of nonhealing. The scenario does not state what type of wound care the patient was receiving. In the presence of PAD, the type of wound care does not matter as much until the PVD is addressed first. Without adequate blood flow to a wound, the wound will not have the nutrients and cells necessary to heal. Bader MS. Diabetic foot infection. Am Fam Physician. 2008;78(1):71-79.

3. A 56-year-old male with hypertension, diabetes presents to the emergency department with redness and purulence draining from his right great toe. His heart rate is 110

beats/min, his blood pressure is 100/80 mm Hg, and his temperature is 38.5° C. Laboratory tests show a white blood count of $14,000/\mu$ L. An x-ray of his foot is performed, which demonstrates gas in his right great toe. What is the best course of treatment?

- A. Outpatient treatment with oral antibiotics and dressing changes.
- B. Inpatient treatment with intravenous antibiotics alone.
- C. Inpatient treatment with intravenous antibiotics and dressing changes.
- D. Inpatient treatment with intravenous antibiotics and emergent surgical debridement .

Answer

D. Inpatient treatment with intravenous antibiotics and emergent surgical debridement.

Discussion: This patient is septic from a diabetic wound infection. He is febrile and tachy-cardic, and he has relative hypotension. This type of patient should not be treated as an outpatient. The correct treatment course involves admission, broad-spectrum antibiotics, and source control with surgical debridement. Gas in the tissues on x-ray in the setting of an infection is an ominous sign and should always prompt surgical debridement. Hence, admission with intravenous antibiotics alone or admission with intravenous and dressing changes alone are inadequate management. Hingorani A, LaMuraglia GM, Henke P, et al. The management of diabetic foot: a clinical practice guideline by the Society for Vascular Surgery in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine. J Vasc Surg. 2016;63(2S):3S-21S.

- 4. A 72-year-old female with diabetes has a chronic right heel ulcer that persists despite local wound care. A magnetic resonance imaging scan demonstrates osteomyelitis of the calcaneus, and a metal probe inserted into the ulcer can "probe to bone." She has a palpable ipsilateral dorsalis pedis pulse. Which operation would be most appropriate?
- A. Transmetatarsal amputation.
- B. Below-knee amputation.
- C. Local calcaneus excision with full-thickness skin flap.
- D. Above-knee amputation.

Answer

B. Below-knee amputation

Discussion: Osteomyelitis of the calcaneus is a difficult condition to treat. The calcaneus is crucial for weight bearing and skin flaps are unlikely to heal in this location. The below-knee amputation will remove the infection and still provide the longest amputation

stump to decrease the work associated with ambulation. As such, an above-knee amputation is too aggressive. A transmetatarsal amputation will not remove the infected bone. Lower extremity amputation: techniques and results. In: Cronenwett JL, Johnson KW, eds. Rutherford's Vascular Surgery. 8th ed. Philadelphia, PA: Elsevier; 2014. UpToDate (Publisher Subscription Required): Lower extremity amputation.

- 5. A 70-year-old male with poorly controlled diabetes and no smoking history presents to the clinic for an ischemic toe ulcer. What level of infrainguinal arterial occlusive disease is he most at risk for developing based on this history?
- A. Superficial femoral artery disease
- B. Deep femoral artery disease
- C. Iliac artery disease
- D. Aortic disease
- E. Tibial artery disease

Answer

E. Tibial artery disease

Discussion: Risk factors for peripheral arterial disease include age, male gender, hypertension, diabetes, smoking, dyslipidemia, family history, and homocysteinemia. Individuals who are heavy smokers tend to develop superficial femoral artery disease, whereas those with diabetes and renal failure tend to develop infrapopliteal disease. The tibial artery is the only infrapopliteal vessel listed as an answer option. All other options are suprapopliteal.

- 6. A 75-year-old male with a history of migraines, diabetes, hyperlipidemia, hypertension, triple coronary bypass (CABG x3), recurrent cellulitis, glaucoma and benign prostatic hyperplasia (BPH) presents to your clinic due to pain. He has smoked approximately 2 packs per day for over 30 years. Despite smoking cessation counseling, he continues to smoke 1-2 cigarettes a day. He reports pain with walking for the past several years. After walking one city block, his calves begin to hurt. He intermittently develops toe sores and wounds that can take 1-2 months to heal. If the wound is larger, it can take 3-4 months to heal. What risk factors does this patient have that contribute to the most likely disease process?
- A. Migraines
- B. BPH
- C. Glaucoma

- D. Smoking
- E. Cellulitis

D. Smoking.

Discussion: Smoking contributes to worsening peripheral arterial disease (PAD). Peripheral arterial disease results when the arteries narrow due to plaque accumulation which prevents or decreases the amount of oxygen-rich blood reaching your legs. The chemicals in tobacco products increase the rate of plaque formation in arterial walls thereby narrowing the arterial lumen. Smoking also constricts blood vessels and causes blood to clot, ultimately putting PAD patients at a higher risk of further complications such as heart attacks, limb amputations, and death. Cellulitis and other wound complications can be an effect of peripheral arterial disease, but is not the cause of peripheral disease. The other options of migraines, BPH, glaucoma have no known associations with PAD.

- 7. A 74-year-old female with a past medical history of IDDM with ESRD on dialysis, hypercholesterolemia, hypertension, and previous MI is referred to you for evaluation of crampy calf pain for the last 2 months with onset upon physical exertion. It has since progressed to pain at rest. You perform ABIs which reveal critical limb ischemia. Which of the following will you most likely observe when performing preoperative imaging of this patient prior to infrapopliteal bypass?
- A. Critical occlusion of only the medial and lateral geniculate arteries.
- B. Loss of enhancement in the region of the popliteal artery but reconstitution showing good distal vessel runoff.
- C. Enhancement in the peroneal artery contributing to pedal circulation.
- D. Complete lack of runoff past the region of the Hunter's canal.
- E. Full enhancement in the posterior tibial artery ending in the region of the medial malleolus.

Answer

C. Enhancement in the peroneal artery contributing to pedal circulation.

Discussion: Because of its location in the deep compartments of the lower leg, the peroneal artery becomes the most likely artery to remain patent in diabetic patients with vascular disease. Therefore, when performing preoperative imaging to plan an infrapopliteal bypass, a more common target site for anastomosis is the peroneal artery. It is important to remember that it is necessary for the peroneal artery to have patency to the anterior or posterior tibial artery to perfuse the entire foot. Choice B is incorrect because this is

more of a presentation in a patient with acute occlusion of only the popliteal artery, as in a posterior knee dislocation. Patients with vascular disease tend to have more distal vessels affected before proximal. Choice A is incorrect because the geniculate arteries themselves are collaterals. Although there is a chance that these will be affected by vasculopathy, it is unlikely that only they will be affected. D is a representation of imaging in a patient with femoral artery disease. Choice E is a scenario that would rarely ever occur without proximal tibial artery being occluded.

Introduction



🅊 Tip

The term peripheral arterial disease (PAD) is often used interchangeably with peripheral vascular disease (PVD) when referring to arterial blockages.

Chronic limb-threatening ischemia (CLTI) is defined as the presence of peripheral arterial disease (PAD) with associated ischemic rest pain, gangrene, or lower limb non-healing ulceration for greater than 2 weeks duration

The prevalence of PAD is approximately 6% in patients aged 40 years or older in the United States. Amongst patients with PAD, 11% of patients (approximately 2 million individuals) will present with CLTI within their lifetime. (Conte et al. 2019c)

CLTI has an increased risk of amputation, cardiovascular morbidity, and mortality.

Etiology

CLTI is a severe manifestation of peripheral arterial disease (PAD), a disease caused by the build-up of atherosclerotic plaque in the arteries. The pattern of occlusive disease can be found in one or a combination of locations including the aortoiliac or infrainguinal vessels (i.e. femoral, popliteal, and tibial arteries) with varying lesion lengths. The superficial femoral artery (SFA) is the most common location where lesions occur. Amongst diabetics, it is typical to find heavier disease burdens more distally, such as the infrapopliteal, tibial vessels (i.e. the tibial trunk, anterior tibial, posterior tibial, and peroneal arteries).

Tip

Here is an image that demonstrates the various peripheral vessels that can be diseased and occluded in a patient with CLTI.

Risk factors for CLTI are the same as for PAD and include advanced age, race (with nonhispanic blacks having the highest risk), male gender, smoking, diabetes mellitus, dyslipidemia, hypertension, and renal insufficiency. It is important to note that any other etiologies including venous, traumatic, embolic, or nonatherosclerotic disease processes are NOT considered a part of chronic limb-threatening ischemia.

Note

For a more comprehensive review of lower limb arterial anatomy and PAD risk factors, please see ?@sec-claudication.

Diagnostics and Imaging

The diagnosis of CLTI requires a prior diagnosis of PAD in association with ischemic rest pain or tissue loss. Pain should be present for more than two weeks and associated with at least one abnormal hemodynamic parameter.

A complete evaluation of patients with CLTI should include:

- 1. A good patient history
- 2. Physical examination
- 3. Noninvasive hemodynamic tests
- Ankle-Brachial Index (ABI) and Pulse Volume Recording (PVR)
- Arterial Doppler Ultrasound
- 4. Angiography
- CTA (abd + pelvis + lower extremity with triple vessel runoff)
- If necessary, invasive angiography

🕊 Tip

For a comprehensive description of the arterial patient history and physical exam, please see ?@sec-patienthistoryandphysicalexaminationofthearterialsystem.

Patient History

- Patients may state that they have persistent pain in their legs, even at rest.
- Rest pain and nonhealing wounds are highly indicative of CLTI.
- Patients may also share experiencing rest pain in bed relieved by dangling their feet off the side of their bed. By dangling their feet over the side of the bed, gravity increases blood flow through diseased and collateral vessels thereby reducing ischemia and pain.

Physical Exam

Common physical exam findings on the lower extremities include:

- Cyanosis
- Cool Extremities
- Hair loss
- Dry skin
- Muscle atrophy
- Dystrophic toenails (thick toenails)
- Capillary refill > 5 seconds
- Non-palpable distal pulses
- Non-healing wounds

Non-invasive Hemodynamic Testing

In terms of non-invasive testing, as stated above, a combination of ABI and PVR is the gold standard in establishing a diagnosis. As a review, the ABI is a non-invasive test that compares blood pressure in the ankle with blood pressure in the arm. ABIs are calculated by dividing the systolic ankle pressure by the highest systolic arm/brachial pressure (between the two arms).

An ABI value: - X > 1.2 is non-compressible, calcification of arteries

- 0.9 < x < 1.2 is normal
- 0.7 < x < 0.9 reflects a mild disease burden
- 0.4 < x < 0.7 reflects a moderate disease burden
- x < 0.4 reflects a severe disease burden

Here is an example of ABI/PVR recordings. Patient A (left) shows a normal ABI/PVR recording, while patient B (right) shows an abnormal ABI with diminished waveforms.

Arterial Doppler ultrasound can also be utilized to evaluate blood flow and detect arterial stenosis or occlusion.

For a definitive CLTI diagnosis, pain should be present for more than two weeks and associated with at least one of the following abnormal hemodynamic parametersI:

- 1. Ankle-brachial index (ABI) < 0.5.
- 2. Absolute toe pressure (TP) < 30 mmHg.
- 3. Absolute Ankle Pressure (AP) < 50 mmHg.
- 4. Transcutaneous pressure of oxygen (TcPO2) < 30 mmHg.
- 5. Flat or minimal pulsatile volume recording (PVR) waveforms.

Angiography

CT Angiography (CTA) with tri-vessel runoff (i.e. anterior tibial, posterior tibial, peroneal arteries) of lower extremities can help identify the location(s) of disease. However, this investigation requires a contrast load and should be carefully considered for those patients with renal disease.

Lastly, diagnostic angiography is an invasive imaging technique that provides detailed information about the arterial anatomy and helps identify specific sites of blockage or stenosis. Diagnostic angiography involves the patient undergoing a femoral artery puncture to access the arterial system. Next, a series of catheters and wires are used to deliver contrast. Fluoroscopic imaging of the lower limb reveals vessel anatomy and identifies areas of stenotic disease. As arterial access is already achieved, immediate intervention may be considered including angioplasty (i.e. ballooning of stenosis) or stenting.

Classification Systems



🕊 Tip

Below, we will introduce and summarize the most common classification systems. For a high-yield summary of all classification systems, please see this paper. (Hardman et al. 2014)

The Rutherford Classification and Fontaine Grading

- The Rutherford Classification is the mainstay of diagnosing the severity of CLTI, however, Fontaine grading can also be used.
- These grading systems are based on symptom presentation. Patients with CLTI usually present with Fontaine Grade 2 or higher or Rutherford Category 3 or higher.
- Minor tissue loss is considered defined as non-healing ulcers with focal gangrene that do not exceed the foot digit.
- Major tissue loss extends beyond the transmetatarsal level with the functional foot no longer salvageable. Patients with major tissue loss most often receive amputations.

Fontaine Grade	Rutherford Category	Clinical Description
0	0	Asymptomatic
	1	Mild claudication
1	2	Moderate claudication
	3	Severe claudication
2	4	Ischemic rest pain
3	5	Minor tissue loss
	6	Major tissue loss

The Wound, Ischemia, and foot Infection (WIfI) classification

- The Wound, Ischemia, and foot Infection (WIfI) classification was proposed in 2014 as an integrated lower extremity wound classification and prognositization system. The WIFI classification was inspired by the tumor, node, metastasis (TNM) staging system for
- Incorporating the presence and severity of lower extremity wounds, ischemia, and infection, the classification system informs providers of the risk of lower extremity amputation and potential benefit of revascularization.
- For the limb in question, a severity grade of 0 to 3 is assigned to the severity and extent of wound, ischemia and foot infection, respectively, where 0 presentes none, 1 mild, 2, moderate and 3 severe disease.

- Using the table here, the combination of these three scores assigns the limb in question to one of four threatened limb clinical stages.
- While an oversimplication, the premise of the WIfI system is that the risk of amputation increases as the disease burden of the limb in question progresses from WIfI stage 1 (very low risk) to tage 4 (high risk).

Wound (ulcer or gangrene)	Ischemia (Toe Pressure/TcPO2 and ABI)	foot Infection
0: No ulcer or no gangrene	0: TP $>= 60 \text{ mmHg}$, ABI $>= 0.80$	0: Not infected
1: Small ulcer and no gangrene	1: $TP = 40-59 \text{ mmHg}$, ABI 0.60-0.79	1: Mild (= < 2 cm cellulitis)
2: Deep ulcer or	2: TP 30-39 mmHg,	2: Moderate (> 2 cm cellulitis or
gangrene limited to toes	ABI 0.40-0.59	involving structures deeper than skin and subcutaneous tissues)
3: Extensive ulcer or extensive gangrene	3: TP $< 30 \text{ mmHg}$, ABI $= < 0.39$	3: Severe (systemic response / SIRS)



The SVS Guidelines Mobile App has an easy-to-use WIfI Calculator. - Apple App Store - Android Google Play

The Global Limb Anatomic Staging System (GLASS) Classification

- The Global Limb Anatomic Staging System (GLASS) is a framework to estimate the success of lower limb revascularization based on the extent and the distribution of the atherosclerotic lesion. By categorizing the patient's lesion into one of three GLASS stages, the surgeon can predict the likelihood of immediate technical failure and one-year limb-based patency (LBP) following endovascular intervention. This information is extremely informative when planning a surgical intervention.
- To perform GLASS staging, the surgeon must obtain high quality imaging of the lower extremity to score the length and burden of disease femoropopliteal segment (FP) and infrapopliteal segment (IP).

Scoring System below (Wijnand et al. 2021)

Femoro-Popliteal (FP) Grading	
0	Mild or no significant (<50%) disease
1	Total length SFA disease $<1/3$ (<10 cm); may include single focal
	CTO (<5 cm) as long as not flush occlusion; popliteal artery with mild or no significant disease
2	Total length SFA disease 1/3-2/3 (10-20 cm); may include CTO
	totaling <1/3 (10 cm) but not flush occlusion; focal popliteal
	artery stenosis <2 cm, not involving trifurcation
3	Total length SFA disease $>2/3$ (>20 cm) length; may include any
	flush occlusion $<20~\mathrm{cm}$ or non-flush CTO 10–20 cm long; short
	popliteal stenosis 2–5 cm, not involving trifurcation
4	Total length SFA occlusion $>$ 20 cm; popliteal disease $>$ 5 cm or
	extending into trifurcation; any popliteal CTO

Infra-Popliteal (IP) Grading	
0	Mild or no significant (<50%) disease
1	Focal stenosis <3 cm not including TP trunk
2	Total length of target artery disease <1/3 (<10 cm); single focal CTO (<3 cm not including TP trunk or target artery origin)
3	Total length of target artery disease 1/3–2/3 (10–20 cm); CTO 3–10 cm (may include target artery origin, but not TP trunk)
4	Total length of target artery disease $>2/3$ length; CTO $>1/3$ (>10 cm) of length (may include target artery origin); any CTO of TP trunk

Trifurcation involvement is defined by disease that include the origin of either the anteriortibial or tibioperoneal trunk.

The combined scores of the FP and IP segments can then be used to determine the GLASS stage using the table below:

		FP Grade				
		0	1	2	3	4
IP Grade	0	NA	Ι	Ι	II	III
	1	I	Ι	II	II	III
	2	I	II	II	II	III
	3	II	II	II	III	III

	FP Grade				
4	III	III	III	III	III

Stage I: Average Complexity Disease: immediate technical failure <10% AND >70% 12-month LBP. Stage II: Intermediate Complexity Disease: immediate technical failure <20% AND 12-month LBP 50-70%. Stage III: High Complexity Disease: immediate technical failure >20%; OR <50% 12-month LBP.

Note

For a more complete explanation of the The Global Limb Anatomic Staging System (GLASS), please see here.

Tip

The SVS Guidelines Mobile App has an easy-to-use GLASS Calculator. - Apple App Store - Android Google Play

The Trans-Atlantic Inter-Society Concensus (TASC classification)

In 2000, a group of fourteen societies from North America and Europe representing a variety of disciplines including vascular surgery, internal medicine, interventional radiology, and cardiology convened to create a consensus in the classification and treatment of patients with peripheral artery disease (i.e. intermittend claudication, acute limb ischemia, and chronic limb ischemia).

The Trans-Atlantic Inter-Society Consensus (TASC) document focuses on individual arterial segments and classifies lesions into one of four large groups (TASC A through TASC D) according to the pattern of disease. Recommendations are then offered according to TASC Classification.

Note

Please see these depictions of the TASC classification system: Image 1 & Image 2.

- TASC A lesions are those that should have excellent results from endovascular management alone.
- TASC B lesions are those that should have good results from endovascular management and endovascular interventions should be the first option offered to the patient.

- TASC C lesions are those that have better long-term results with open surgical management. Endovascular techniques should be reserved for patients who are poor surgical candidates.
- TASC D lesions are those that should be treated by open surgery.

Important caveats: - Developments in endovascular techniques and devices since the introduction of the TASC document have increased the role of endovascular surgery in treating TASC C and TASC D lesions.

 Many patients with CLTI present with multisegment disease thereby complicating the clinical decision-making usefulness of TASC recommendations. TASC is more reasonably useful for assessing lesion-specific device performance.

The consensus was updated in 2007 (TASC II) and included representatives from Australia, South Africa, and Japan.

Treatment

Treatment of CLTI includes conservative management and invasive intervention. Lifestyle modification is recommended and highly encouraged and includes smoking cessation, regular exercise, and a healthy diet to reduce cardiovascular risk factors.

Conservative management

Pharmacotherapy with antiplatelet agents and statins plays a large role in helping prevent the progression of CLTI. Antiplatelet agents – such as aspirin or clopidogrel, reduce the risk of blood clots, while statins control cholesterol levels and reduce the progression of atherosclerosis. In some instances, vasodilators, such as cilostazol, can be utilized to improve blood flow and relieve symptoms, especially in those individuals experiencing ischemic rest pain.

Invasive management

The mainstay of treatment modality in CLTI is revascularization with endovascular therapy, surgical bypass, or a combination of both.


Endovascular revascularization is the least invasive approach to opening narrowed or blocked arteries. Using real-time contrast-enhanced images, the surgeon can perform angioplasty (i.e. ballooning of stenosis) stenting, and/or atherectomy (i.e. device mediated removal of the intimal layer and atherosclerotic material).

A surgical bypass, or extra-anatomic bypass, is the surgical creation of a new path for blood around the area of vessel stenosis/occlusion. Conduits for bypass can be artificial (e.g. PTFE or dacron grafts) or natural (e.g. autogenous or cadaveric saphenous vein grafts). In severe cases where limb salvage is not possible or limb-threatening infection persists even after attempts at revascularization, amputation may be necessary to preserve overall health and quality of life.

In addition to conservative management and surgical intervention, appropriate wound care is an important aspect of CLTI treatment. Surgical wound debridement, infection control, offloading techniques, and daily specialized wound care are integral in promoting healing and preventing amputation.

It is important to note that CLTI is a chronic disease, and patients may require multiple interventions, be on lifelong aspirin/statin therapy, and require close follow-up with routine imaging surveillance for the entirety of their lifetime. Moreover, vascular surgeons work with a team of physicians from other specialties – including primary care, cardiology, podiatry, nephrology, and endocrinology – in order to ensure that patients with CLTI are appropriately optimized given their respective comorbidities to improve overall outcomes. This integrative physician network to help manage CLTI patients is important as patency rates of stents and bypasses can be severely limited by a patient's underlying disease burdens from uncontrolled hypertension, diabetes, renal insufficiency, and tobacco abuse.

Below is a sample algorithm for treatment management of a patient with PAD that progresses to CLTI. Candidacy for revascularization indicates that the patient is able to tolerate anesthesia and has suitable anatomy - including possible targets for bypasses or the ability to traverse lesions endovascularly.



Here is an image demonstrating angioplasty, atherectomy, and placement of stents.

Here is an image demonstrating surgical or extra-anatomic bypass.

Outcomes

- Patients with CLTI have variable outcomes with regard to limb loss and overall mortality.
- Approximately 90% of all CLTI patients undergo a revascularization procedure (bypass or endovascular intervention) in their lifetime. (Conte et al. 2019d)
- For CLTI patients who do not undergo any revascularization, the natural history includes a 1-year major amputation and 1-year mortality rate of 22%. (Conte et al. 2019d)
- When looking at overall 1-year outcomes for all CLTI patients, 1-year amputation rates are at 15-20% and 1-year mortality is at 15-40%. (Conte et al. 2019d)

- Less than half of all CLTI patients (45%) will have both limbs after 1-year from initial time of diagnosis. These rates are even higher amongst the subset of CLTI patients who have diabetes. (Conte et al. 2019d)
- As the diagnosis of CLTI carries a high risk of mortality and morbidity even with vascular intervention to improve blood flow, there is a strong emphasis on preventing the progression of PAD to CLTI. (Conte et al. 2019d)

Surveillance

Medical Treatment Surveillance

- Patients with CLTI have variable outcomes with regard to limb loss and overall mortality.
- After revascularization procedures, Patients remain on dual antiplatelet therapy (DAPT) for 1-6 months.
- Through DAPT has been show to have increased bleeding risk, it may result in increased survival and reduced risk of amputation. (Conte et al. 2019d)
- There is not a clear benefit to the use of of cilostazol, a PDE inhibitor, after endovascular interventions for CLTI. (Conte et al. 2019d)

Endovascular Surveillance

- Early failure of endovascular procedures is common. As such, it is recommend that patients undergo duplex ultrasound surveillance and duplex guided interventions to prompt long-term patency. However, there are no set guidelines for how long and how frequently to perform surveillance.
- Recommended methods of surveillance include clinic visits, ankle-brachial indexes (ABIs), and duplex scans.
- Surveillance intervals can range from 3-6 months.
- Ultimately, there is no strong evidence of benefit of routine duplex surveillance after endovascular intervention of CLTI; however, patients with the characteristics below may benefit from duplex surveillance. (Conte et al. 2019d)

Patient Risk Factors that may Prompt Duplex Surveillance

Multiple failed angioplasties and failed bypasses Severe ischemia Unresolved tissue loss Poor Inflow

Vein and Prosthetic Bypass Surveillance

Surveillance of vein grafts after bypass surgery may solely be clinical and may include the vascular laboratory as well. The SVS recommends a 2 year surveillance program postoperatively to detect new symptoms, monitor changes in ABIs, and monitor for stenotic lesions within graft or at surveillance sites.

- Duplex surveillance and intervention based on duplex findings lead to better patency rates of the vein graft.
- Though there are no set guidelines for frequency of surveillance. Many studies in the literature often perform surveillance 30 days postoperatively and at 3-6 months intervals.
- Ultimately, though duplex surveillance does result in better patency of vein grafts, there
 is no clear evidence that duplex surveillance for CLTI results in clinical benefits for
 patients.

Prosthetic bypasses are known to fail more frequently and more early compared to vein bypasses. Similar to vein bypassess, the SVS does not have hard recommendations for when to perform duplex surveillance.

- Many studies perform duplex at 30 days postoperatively and at 6 month intervals.
- Despite this, the duplex itself does not always always reliably detect intervenable lesions as it does in vein bypasses.

Teaching Case

Scenario

A 73-year-old male with a significant smoking history and medical history including 4 vessel coronary bypass 10 years ago, hypertension, hyperlipidemia, and diabetes presents to clinic for pain in his lower extremities. He states that it is difficult for him to walk 3 blocks without having to stop. He locates the pain in his calves bilaterally. He endorses he has experienced this pain for over 5 years. He mentions that he previously had a wound that took almost 3 months to heal. He denies presence of new sores or wounds. He had a right lower extremity angiogram 2 months ago that failed to recanalize the superficial femoral artery (SFA).

Exam

HEENT: No prior neck incisions, no lymphadenopathy.

Cardiac: Regular rate and rhythm (RRR), healed stereotomy scar.

Pulmonary: Clear to auscultation bilaterally, no increased work of breathing.

Abdominal: Soft, non-distended (ND), non-tender (NT)

Vascular/Extremities: Non-palpable lower extremity pulses and what appears to be a wound that is starting to form on the distal big toe of the right foot. He does have 2+ femoral pulses. On Doppler exam, he has monophasic dorsalis pedis (DP) and posterior tibial (PT) signals in his right lower extremity.

Imaging

Duplex Ultrasound: On duplex ultrasound, he has a long segment occlusion of the superficial femoral artery (SFA) and on CTA he has two-vessel run off to the right lower extremity with well-formed collateralization.

Discussion Points

N.B. There is no prepared answers for the questions below.

However, we feel this chapter contains all the necessary information to answer the questions. If not, please let us know!

- 1. Please explain the pathophysiology of the patient's lower extremity pain and non-healing wounds.
- 2. Please list the patient's risk factors for Chronic Limb Threatening ischemia.
- 3. Explain why there is well-formed collateralization that was seen on CTA.
- 4. What next steps should be pursued to pursue treatment for the patient? Should the patient stop smoking?
- 5. What surgical treatment could be offered to this patient?
- 6. What additional imaging would help with surgical planning for the patient?

Key Articles

- Beckman JA, Schneider PA, Conte MS. Advances in Revascularization for Peripheral Artery Disease: Revascularization in PAD. Circ Res. 2021;128(12):1885-1912. doi:10.1161/CIRCRESAHA.121.318261.(Beckman, Schneider, and Conte 2021)
- 2. Berchiolli R, Bertagna G, Adami D, Canovaro F, Torri L, Troisi N. Chronic Limb-Threatening Ischemia and the Need for Revascularization. JCM. 2023;12(7):2682. doi:10.3390/jcm12072682.(Berchiolli et al. 2023)
- 3. Blanchette, Virginie, Malindu E. Fernando, Laura Shin, Vincent L. Rowe, Kenneth R. Ziegler, and David G. Armstrong. Evolution of WIfI: Expansion of WIfI Notation After Intervention. The International Journal of Lower Extremity Wounds, 2022. doi:10.1177/15347346221122860. (Blanchette et al. 2022)
- 4. Conte MS, Bradbury AW, Kolh P, et al. Global vascular guidelines on the management of chronic limb-threatening ischemia. Journal of Vascular Surgery. 2019;69(6):3S-125S.e40. doi:10.1016/j.jvs.2019.02.016.(Conte et al. 2019e)
- 5. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the Management of Patients With Lower Extremity Peripheral Artery Disease: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. Circulation. 2017;135(12). doi:10.1161/CIR.000000000000000470.(Gerhard-Herman et al. 2017b)
- Mills JL, Conte MS, Armstrong DG, et al. The Society for Vascular Surgery Lower Extremity Threatened Limb Classification System: Risk stratification based on Wound, Ischemia, and foot Infection (WIfI). Journal of Vascular Surgery. 2014;59(1):220-234.e2. doi:10.1016/j.jvs.2013.08.003.(Mills et al. 2014)
- 7. Wijnand JGJ, Zarkowsky D, Wu B, van Haelst STW, Vonken EPA, Sorrentino TA, Pallister Z, Chung J, Mills JL, Teraa M, Verhaar MC, de Borst GJ, Conte MS. The Global Limb Anatomic Staging System (GLASS) for CLTI: Improving Inter-Observer Agreement. J Clin Med. 2021 Aug 4;10(16):3454. doi: 10.3390/jcm10163454. PMID: 34441757; PMCID: PMC8396876.(Wijnand et al. 2021)

Additional Resources

Audible Bleeding Content

• Audible Bleeding Exam Prep: CLTI Chapter

Gore Combat Manual

The Gore Medical Vascular and Endovascular Surgery Combat Manual is an informative and entertaining read intended as a vascular surgery crash course for medical students, residents, and fellows alike. Highly accessible with a thoughtfully determined level of detail, but lacking in learning activities (e.g. questions, videos, etc.), this resource is a wonderful complement to the APDVS eBook.



Please see pages 85-98.

Operative Footage

Developed by the Debakey Institute for Cardiovascular Education & Training at Houston Methodist. *YouTube account required as video content is age-restricted.* Please create and/or log in to your YouTube account to have access to the videos.

External Iliac Artery Balloon Angioplasty and Stenting

Common Femoral Endarterectomy

Left Femoral to Peroneal Bypass by Reversed Ipsilateral Great Saphenous Vein (Surgical or Extra-anatomic Bypass)

Left Common Femoral Artery to Anterior Tibial Artery Bypass using PTFE Graft (Surgical or Extra-anatomic Bypass)

Below Knee Amputation (BKA)

Above Knee Amputation (AKA)

Miscellaneous Topics

Basics of Endovascular Surgery (Endo Basics)

Jeremy Zack, Nitin Jethmalani, MD, Babak Abai, MD, FACS.

Note

By the end of this chapter, students will:

- Be able to define endovascular surgery.
- Be able to elaborate on the key principles of arterial access, wires, catheters, and sheaths.
- Be able to outline the steps and associated risks of common endovascular surgeries.
- Be able to outline new developments in endovascular surgery.

Key Facts

- 1. Endovascular surgery is a minimally invasive approach to treat a variety of vascular pathologies, including peripheral artery disease, aortic aneurysms, venous diseases, and others by deploying baloons, stents, and or endografts.
- 2. The common femoral artery (CFA) is the most common site of access for endovascular surgery due to its larger size and lower risk of complications relative to other sites of vascular access
- 3. Using the analogy of a train on a track passing through a tunnel, a catheter can be thought of as the train, a wire as the train track, and a sheath as the surrounding tunnel.
- 4. Wires are used to guide devices such as catheters and sheaths into a vessel.
- 5. Catheters are flexible, hollow tubes that are used in both the diagnosis and treatment of vascular disease. In an analogy of a train track, the catheter can be thought of as the train, the track is the wire, and the surrounding tunnel is the sheath.
- 6. Catheters are measured in French (1F = 0.33 mm) and the diameter corresponds to the outer diameter of the tube.
- 7. Sheaths are hollow tubes that provide conduit between outside the body and into the vessel, the "tunnel" in the train track analogy. Sheaths are used to advance

- and exchange catheters, wires, stents, balloons, and other devices.
- 8. Sheaths are measured in French (1F = 0.33 mm) and the diameter corresponds to the inner diameter of the tube.
- 9. A 6F catheter will pass through a 6F sheath.
- 10. Endovascular surgery is not without risks and complications. Complications include the formation of hematoma, pseudoaneurysm, retroperitoneal hematoma, fistula, dissection, thrombosis and embolism, and atheroembolism. Other complications include wire perforation vessel rupture.

• How We Suggest Using the Pre/Post Questions

The pre/post questions are listed below. They are all multiple choice questions with a single right answer. To best guide your learning, we have hidden the answers in a collapsible menu. Before reading the chapter, we suggest giving the questions a try, noting your answers on a notepad. After reading the chapter, return to the questions, reevaluate your answers, and then open the collapsible menu to read the correct answer and discussion. Do not fret if you have difficulty answering the questions before reading the chapter! By the end of the chapter, we are certain you will have covered the knowledge necessary to answer the questions. There will be a teaching case at the end of the chapter. This is another opportunity to exercise your new knowledge!

Pre/Post Questions

Case Based Questions

- 1. A 77-year-old female with a past medical history of CAD, CHF with an EF of 14%, COPD, and a prior total thyroidectomy comes in for evaluation of carotid stenosis. She had a TIA with a right-sided facial droop two weeks ago and was briefly admitted to this hospital. She had a carotid US and was found to have an 85% left carotid stenosis. Which of the following treatment modalities would be best suited for the patient?
- A. Open carotid endarterectomy
- B. Transcarotid Artery Revascularization (TCAR)
- C. No intervention because of her age
- D. No intervention because of her medical comorbidities

B. Transcarotid Artery Revascularization (TCAR)

Discussion: Considering the patient has a symptomatic carotid artery lesion, her disease warrants treatment. This rules out answers C and D. TCAR is a safe, minimally invasive surgical technique used to treat carotid artery disease. TCAR and other endovascular procedures are often a safer option in the elderly, medically comorbid patients, especially those who have a higher risk of complications (such as those with prior neck surgery). In addition, TCAR can be performed under local anesthesia with sedation. This makes TCAR a safer option relative to open CEA, making B a better answer than C.

- 2. A 66-year-old male presents to the emergency room one week after a percutaneous coronary intervention via right groin access. He reports that over the last few days, he has noticed a bulge in his groin and when he touches it, it is painful and has a pulsatile nature. A complete exam reveals a pulsatile, tender mass in the right groin with no skin changes, but with a bruit on auscultation. Which of the following complications is NOT associated with this patient's diagnosis?
- A. Embolization to the lower extremity
- B. Rupture of the mass
- C. True femoral artery aneurysm
- D. Compression of the femoral vein
- E. Gangrene of the toes

Answer

C. True femoral artery aneurysm

Discussion: In the post-intervention patient who presents with a pulsatile tender mass at the intervention site with a bruit on auscultation, the most likely diagnosis is a femoral artery pseudoaneurysm. Risks of an untreated PSA include embolic events, distal ischemia including gangrene, rupture of the PSA, skin necrosis, and compression of surrounding structures including the femoral vein and nerve. Because a PSA is an outpouching of only 1 or 2 of the layers of the arterial wall, a true aneurysm (outpouching of all three layers of the artery, intima, media, and adventitia) is not associated with a femoral PSA.

3. A 42-year-old patient with a history of an arterial thrombus four years ago presents to the ER with severe right lower extremity pain. A CTA is obtained showing no atherosclerotic disease but with acute thrombus along the distal SFA and popliteal artery. You take the patient to the operating room for a lower extremity angiogram. Which catheter would be most useful to help break up the clot over time?

- A. Penumbra thrombectomy catheter
- B. Cragg Mcnamara Catheter
- C. Fogarty Catheter
- D. Peripherally inserted central venous catheter

B. Cragg Mcnamara Catheter

Discussion: In this case, the patient has an acute thrombus in her lower extremity arteries. Although there are many treatment options for this disease process depending on disease burden and characteristics, the question specifically asks about a catheter that can be used to break up the clot over time. The Cragg Macnamara catheter is a thrombolysis catheter that remains in place infusing "clot buster" medication (tPA) over a period of hours to days to dissolve the clot. A Penumbra thrombectomy catheter is used to suction blood clots out of a vessel. A Fogarty catheter can be used to manually withdraw thrombus from a vessel. A PICC line is used for long-term infusions to deliver medications to the central venous system and is not used intra-operatively for procedures.

- 4. A 70-year-old male is brought to the operating room for an EVAR procedure for his 6 cm infrarenal AAA. You have chosen a graft to use and the nurse is asking if there are any wires that you would like opened to deliver the device into the patient's aorta. Which of the following wires would be most useful for this task?
- A. 0.018" flexible Terumo wire
- B. 0.035" Amplatz wire
- C. 0.035" floppy Glidewire
- D. 0.014" Pilot wire

Answer

B. 0.035" Amplatz wire

Discussion: Vascular access wires each have a variety of uses determined by their stiffness, diameter, hydrophilicity, and tip shape. In this case, a stiff wire to pass a large aortic endograft over would be the most useful. The larger diameter wire will be stiffer, which rules out options "A" and "D", which would both be used for getting access to smaller vessels. Option C is a floppy glide wire which can be useful for crossing lesions and getting up and over the aortic bifurcation. The 0.035" Amplatz wire is a stiff guidewire that will not buckle under the pressure of passing a large device into the aorta.

- 5. A 63-year-old smoker presents to the clinic with left lower extremity claudication and an ischemic ulcer. He is taken to the operating room for an angiogram via right groin access. Which of the following steps is INCORRECT when obtaining arterial access?
- A. Using anatomic landmarks to confirm cannulation through the inguinal ligament
- B. Using ultrasound to ensure cannulation of the common femoral artery
- C. Using fluoroscopy to ensure puncture at the level of the femoral head
- D. Using the Seldinger technique for first access, making sure not to lose access to the artery during exchanges or wires and catheters.

A. Using anatomic landmarks to confirm cannulation through the inguinal ligament

Discussion: On first access for an angiogram with right groin access, the right common femoral artery should be accessed. The CFA begins below the level of the inguinal ligament, making answer A incorrect. One should avoid going through the inguinal ligament as native fascial planes are disrupted and it can make for challenging upsizing of devices and closure. Following identification of the inguinal ligament; ultrasound, fluoroscopy, and the Seldinger technique should be used as above to safely cannulate the CFA.

Introduction

Endovascular surgery is a minimally invasive approach to treat a variety of vascular pathologies including peripheral artery disease, aortic aneurysms, venous diseases, and more. Endovascular surgery is not limited to vascular surgery as a specialty and is also employed by interventional cardiologists, neurosurgeons, interventional radiologists, and others.

Over the past 30 years, endovascular surgery has dramatically changed the practice of vascular surgeons, decreasing patient morbidity and mortality in the immediate postoperative period and the length of hospital stay.

Understanding endovascular surgery as a medical student or junior resident can be a difficult task due to the abundance of wires, catheters, sheaths, and devices used and orienting onself using fluorscopic images. In this chapter, we will focus on developing a conceptual basis for endovascular surgery.

Relevant Anatomy and Vascular Access

The common femoral artery (CFA) is the most common site of access for endovascular surgery due to its larger size and lower risk of complications relative to other sites of vascular access. Other sites of arterial access include the radial, brachial, and axillary artery in the upper extremity, as well as the popliteal and dorsal pedal arteries in the leg. The further away the vessel is from the aorta, the smaller the vessel lumen, and hence the smaller the diameter of devices that can be utilized in that artery.

The CFA is referred to as the external iliac artery proximal to the inguinal ligament and the superficial femoral artery (SFA) distal to the branching of the profunda femoris artery (PFA). The anatomic relationship of the CFA to the surrounding vessels can be remembered with the mnemonic "NAVeL", femoral nerve, artery, vein, and lymph nodes from lateral to medial. The Femoral Triangle consists of these three vessels and nerves surrounded by the inguinal ligament superiorly, sartorius muscle laterally, and the adductor longus medially.




Figure 1: Femoral triangle anatomy courtesty of S Bhimji, MD.

Correct anatomic positioning for cannulation of the CFA can be achieved by palpating the point of maximum pulsation inferior to the inguinal ligament, a structure identifiable by tracing along the facial plane between the anterior superior iliac spine and pubic symphysis. The following imaging modalities assist with anatomic positioning:

- Fluoroscopic guidance: cannulation of the CFA should be attempted at the level of the middle to inferior femoral head.
- Ultrasound (US) guidance: identification of the PFA ensures cannulation is attempted of the CFA in place of the SFA. At the level where the great saphenous vein joins the femoral vein, US will show one large vessel surrounded by two smaller vessels, which can be remembered as the "Mickey Mouse Sign."



Tip

An in depth review of lower limb anatomy can be found in the APDVS Claudication Chapter or here.

Ultrasound Guided Access

There is emerging evidence to suggest real time identification of anatomic structures with US guidance may decrease the risk of access-related complications. A summary to the steps of US-guided access of the CFA are included below.

Identifying the correct vessel:

- 1. Obtain baseline peripheral pulses.
- 2. Obtain a micropuncture kit, containing a hollow needle, guidewire, and catheter
- 3. Using the US probe, identify any plaque or calcification along the CFA or iliac artery that may change the site of access.
- 4. Identify the CFA, which can be differentiated from the femoral vein by its thick wall, resistance to compression, pulsatility, and relative lateral anatomic positioning. A video identifying the CFA through the "Mickey Mouse Sign" on US is available here.

Obtaining access through the Seldinger technique:

The Seldinger technique help obtain **safe** access to blood vessels. In contrast to previous methods, the Seldinger technique causes minimal damage to the blood vessel thereby allowing the puncture to be closed percutaneously, and avoids the risk of loss of access during the procedure through keeping at least one wire, catheter, or sheath at the site of interest in the body all times.

Steps to the Seldinger technique are as follows:

- 1. Cannulate the CFA by inserting the hollow needle at a 45-90 degree angle to the skin under US guidance, blood will flow out of the needle once inserted into the vessel.
- 2. Insert the guidewire into the vessel. Remove the hollow needle while keeping the guidewire in place, being careful not to allow the wire to withdraw as the needle is removed.
- 3. Insert the catheter over the guidewire.
- 4. Exchange the guidewire that came in the micropuncture kit with desired guidewire to begin the procedure.

Please see this video animation of the Seldinger technique for vascular access.

Closure and Closure Devices

Once the procedure is complete, percutaneous access will need to be closed. If relatively small catheters and sheaths were used during the procedure, generally below 6 French (1F=.33 millimeter), arterial access may be closed with manual compression by applying pressure with two fingers over the site of incision against the femoral head. Alternatively, certain indications, such as use of anticoagulation or large body habitus, may require closure devices even if larger catheters or sheaths were not used during the procedure.

There are several different types of closure devices. Each type can be differentiated by the use of suture or sealant to close the vessel.

A common closure device you are likely to encounter on a vascular surgery service is the **Perclose ProGlide** (Abbott Vascular, Chicago, IL), a suture-mediated closure device. Other closure devices include the **Angioseal** (Terumo, Tokyo, Japan), which deploys a bioabsorbable anchor to the arterial puncture site with surrounding collagen to promote closure, and the **MYNXGRIP** (Cordis, Miami Lakes, FL), which uses a balloon pressed against the site of arterial puncture and an adhesive sealant outside the vessel that dissolves slowly as the puncture site heals. Less commonly used closure devices include the **Angioseal** (Terumo, Tokyo, Japan), which deploys a bioabsorbable anchor to the arterial puncture site with surrounding collagen to promote closure. Other less commonly used closure devices include the **Manta**, **StarClose**, **Vascade**, and **Cardiva**.

A step-by-step guide to deploying a **Perclose device** is shown in this animated video and listed below.

- 1. Exchange the procedural sheath for a guide wire so only the guide wire is left in the vessel.
- 2. Advance the flushed perclose device over the guidewire until the wire exit port is at skin level
- 3. Remove the guidewire so only the perclose device is in the vessel and advance the perclose device until blood flows out of the marker lumen.
- 4. Lift the lever to open the foot of the device.
- 5. Insert the plunger down to insert needles, which will connect the threads of the perclose
- 6. Remove the plunger, which pulls the thread taught, and cut the excess thread Lower the lever to close the foot and retract the device until the guide wire access port is visible.
- 7. Reinsert guidewire and pull both threads taught.

- 8. Use the Snared Knot Pusher or Prostyle Suture Trimmer to advance the knot to the level of skin. 9 Pull the thread taught while removing perclose device.
- 9. Advance the knot over the guidewire and remove the guide wire once the bleeding is controlled.
- 10. Pull both threads tightly and remove the knot advancing device.

Wires, Catheters, and Sheaths



Tip

Using the analogy of a train on a track passing through a tunnel, the catheter can be thought of as the train, the wire as the train track, and the sheath as the surrounding tunnel.

Wires

Wires are the first device inserted into a patient, have the lowest profile, and are variable in properties. Vascular access wires are made of a metal core and an outer wire wrapped around the metal core. They are used to guide devices such as catheters and sheaths into a vessel. Characteristics of wires include:

Diameter

- Wire diameter is described in thousandths of an inch and typically range from 0.010-
- Vascular access is typically obtained with a 0.018" wire and is exchanged for a smaller or larger profile as needed.
- $\bullet\,$ The most common sizes used are 0.035", 0.018" and 0.014".

Stiffness

- Wire stiffness is determined by the metal used as the core of the wire.
- The same wire will become stiffer with a larger diameter.
- Softer wires are used to manipulate catheters into vessels.
- Stiffer wires are needed to push larger devices, such as balloons or endografts.

Coating

- Wires have a hydrophilic or hydrophobic coating.
- Hydrophilic wires have less friction when moving along the vessel wall and track more closely along the vessel lumen, increasing the risk of dissection.
- Hydrophilic wires are preferable in highly tortuous vessels.
- In contrast, hydrophobic wires keep more distance from the vessel wall and are less likely to dissect.

Tip Shape

- The tip shape and consistency of the wire is important.
- Angled tips are used to navigate into side branches of vessels.
- Soft tips are designed to prevent trauma to the vessel wall.
- Weighted tips are used to cross occlusions.
- J-tips are used to prevent wiring of too small of vessels that may lead to trauma of end organs.

Length

- Wires are measured in centimeters (cm) and typically range from 80 300 cm.
- Wires must not only be long enough to reach the target vessel from the point of access, but also need to be long enough to support exchanges outside of the body as needed.
- Care must be taken to avoid too long of a wire for its intended purpose as it is harder to control the end of the wire outside of the body.

Function

Wires can be categorized by function. Please find a list of each category with common examples of specific wires below.

Access wires:

• Short, typically 0.018" wires with a soft, non-directional tip to prevent any trauma to the vessel when entering, and a stiffer body to allow tracking of the introducer and sheath into the vessel.




Figure 2: Commonly used wires, curtosy of B. G. Northcutt, MD, A Shah, MD, Y Sheu, MD, L Carmi, MD.

Guidewires:

- Generally are characterized as pushable, hydrophilic wires with a floppy, curved or angled tips that are designed to track through tortuous vessels without causing blunt trauma to the vessel wall.
- Guidewires are also used for their directionality to cannulate branches.
- Standard Glidewire (Terumo, Tokyo, Japan): can be used as an access or guidewire. Hydrophilic coating, straight or angled tip, sized 0.018" 0.035" in diameter and 150 cm or 260 cm in length.
- Bentson (Cook Medical, Bloomington, IN): flexible wire with a hydrophobic coating and curved tip. Used to access vessels that have a slight curve, or to introduce catheters or sheaths. Comes in 0.035" diameter and 150 cm or 260 cm in length.
- Rosen (Cook medical, Bloomington, IN): stiff hydrophobic wire with J tip. Often used as the first wire exchanged for the access wire. 0.035" long and 80 cm to 260 cm in length.

Rail wires:

- Used as a platform to support catheter and sheath exchanges, as well as to advance devices, such as stents or endografts.
- Rail wires come in a range of stiffness, from medium to very stiff.
- They may have a straight, floppy, or curved tip.

Medium stiffness rail wires:


- Bentson (Cook Medical, Bloomington, IN): flexible wire with a hydrophobic coating and curved tip. Used to access vessels that have a slight curve, or to introduce catheters or sheaths. Comes in 0.035" diameter and 150 cm or 260 cm in length.
- Rosen (Cook medical, Bloomington, IN): stiff hydrophobic wire with J tip. Often used as the first wire exchanged for the access wire. 0.035" long and 80 cm to 260 cm in length.

High stiffness rail wires:

- Amplatz (Boston Scientific, Marlborough, MA): stiff, hydrophobic, straight floppy or J-tip. 0.035" 0.038" diameter and 75 cm 260 cm in length.
- Lunderquist (Cook Medical, Bloomington, IN): one of the stiffest wires, hydrophobic, with floppy or curved tip. 0.035" diameter and 90 cm 300 cm in length. Often used to advance large endografts.

Catheters

Catheters are flexible, hollow tubes that are used in both the diagnosis and treatment of vascular disease.



Catheters vary by size, stiffness, hydrophilicity, radiopacity, and functionality, but can generally be described in two categories, flush catheters and selective catheters.

Flush catheters:

- Contain multiple side holes just proximal to the end of the device to allow for the injection of contrast or other material, such as thrombolytics.
- Omniflush (Angiodynamics, Latham, NY): soft, flexible catheter with a radiopaque tip that is able to administer contrast to visualize vessels. Omniflush device on back table
- Pigtail (Angiodynamics, Latham, NY): Similar to the Omniflush with a looped end (Image YY).

Selective catheters:

- Contain one hole at the distal end of the tube.
- Selective catheters come in many different shapes and sizes.
- Intuitively, the anatomy of the vessel to be tracked will determine the diameter and tip shape.

- When attempting to cross the aortic bifurcation into the contralateral common iliac artery, a catheter with more of a J-tip or "Shepherd's hook" is preferable.
- When attempting to cannulate vessel branches with a more acute angle, a smaller angle catheter tip should be selected.
- Selective catheters can also be used to deliver contrast, but the contrast is only able to leave the catheter through the end of the device.
- Angiograms with selective catheters are more tailored to specific vessels, such as a renal
 or mesenteric artery.
- Selective Glidecath (Terumo, Tokyo, Japan): hydrophilic catheter available in 17 different shapes, but the 45 degree is most common. Comes in 4F or 5F. Offers great trackability and pushability. This image depicts some of the glidecath tip shapes.

Sheaths

Sheaths are hollow tubes that provide a conduit between outer environment (i.e. outside the body) and the vessel. Returning to our train analogy, a sheath is the "tunnel" that the track and train run through.

Sheaths are used to advance and exchange catheters, wires, stents, balloons, and other devices. Sheaths can be divided into short sheaths for access and long sheaths for diagnostics and treatment. Long sheaths are flexible enough to track along different vessels, but are also stiff enough to support the advancement or exchange of devices.

Note

Sheaths are measured in French (1F = 0.33 mm) and the diameter corresponds to the inner diameter of the tube.

As such, a 6F catheter will pass through a 6F sheath.

Key points about sheaths:

- Sheaths are measured based on the inner diameter of the sheath, in contrast to catheters, which are measured based on their outer diameter.
- Sheaths contain a removable introducer, which is used to expand the tract from the outside into the vessel, and provide a smooth transition between the wire and the sheath.
- The side port on sheaths is used to flush the inner device with heparinized saline to prevent thrombus formation and distal embolization. It can also be used to inject contrast for diagnostic purposes.
- Sheaths come in variable diameters, lengths, and tips.
- Some sheaths, such as the **TourGuide** (Medtronic, Minneapolis, MN) are steerable. The operator can deflect the sheath tip up to 180 degrees. This is advantageous when cannulating branch arteries with sharp angles.

It is also helpful to understand what diameter of sheath can be inserted into different points of access as this will change the approach for the procedure. Though there is variation across patients, the table below is a useful starting point to understand what size sheath can be inserted in target access arteries.

Artery	Inner Diameter (mm)	Typical Access Sheath size (F)
Radial	2.3	4-6
Brachial		5-7
Axillary		5-14
Common Femoral	6.6	5-24
Popliteal	4.7	4-6
Pedal	2.1	4

Balloons, Stents, and Endografts

The delivery system for balloons consists of a catheter over the wire with the balloon built into the catheter tip. As such, advancing the balloon into place for angioplasty is no different from advancing a small catheter without a balloon.

Similar to balloons, stents are contained in catheters that are advanced into position over wires. The interventional component (i.e. stent) is near the tip of the catheter.

Both stents and balloons can elute drugs, most commonly *Paclitaxel*, an antiproliferative agent. Paclitaxel is lipophilic and remains active in the vessel wall following the procedure, preventing neointimal hyperplasia and improving patency. The impetus for using a drug-eluting stent or balloon in place of a standard stent or balloon is to decrease the chances of vascular remodeling following the procedure that may lead to restenosis.

In contrast to drug-eluting stents, drug-eluting balloons have the advantage of not leaving hardware in the vessel after intervention. Contrarily, drug-eluting stents remain in the vessel lumen after intervention, but are able to continue providing radial force.

Balloons

Balloons are versatile endovascular tools and can be categorized by the degree to which they assume of the shape of the vessel.

Compliant balloons: - Compliant balloons inflate to a specific volume and *conform to the shape of the vessel.* - For example, the **Fogarty balloon** is a compliant balloon that can be used to remove thrombi in both open and endovascular procedures.

Semi-Compliant balloons: - These balloons have a slightly more rigid shape and therefore conform to a lesser degree to the shape of the container (i.e. vessel). - The Coda balloon (Cook Medical, Bloomington, IN) is semi-compliant and used to temporarily occlude large vessels.

Non-Compliant balloons: - Non-compliant inflate to a predetermined shape and size. - They are commonly used in angioplasty of peripheral and/or visceral stenosed vessels.




Figure 4: Angioplasty with a non-compliant balloon courtesty of J. Heuser, and Wikimedia.

Intraoperatively surgeons will often ask for the pressures of the balloon before using. Non-compliant balloons have a **nominal pressure**, which is the average pressure in atmospheres (atm) required to inflate the balloon to its *predetermined size*, and a **burst pressure**, where 99.9% of balloons will not burst if inflated to this pressure.

Stents

Stents may be covered or uncovered. Covered stents do not allow flow between the lumen of the stent and surrounding tissues, in contrast to uncovered (bare metal) stents which do allow flow through the stent.

Stents can be categorized by the means of deployment.

Self-expanding stents use a recoil mechanism during deployment to achieve a set diameter using stored energy from the packing of the stent.

Key characteristics of **self-expanding stents**:

- Flexible
- Low profile
- Deformable
- Low radial force (lower dissection risk)

Balloon-expandable stents are packaged around a balloon. The surgeon uses the balloon inside the stent to achieve a specific diameter based on the pressure used to inflate the balloon. As the diameter of the stent increases with higher balloon pressure, the length of stent decreases.

Key characteristics of balloon-expandable stents:

- Rigid
- Large profile
- Maintains shape over time
- More accurate deployment
- High radial force (higher risk of dissection)



Highly tortuous vessels often require a **self-expanding stent** because of the increased flexibility needed to prevent kinking or in-stent restenosis.

Heavily calcified vessels often require the increased radial force of balloon-expandable stents to achieve full patency.

Endografts

Endografts are tubes of fabric and interwoven metal used to exclude the diseased segment of an artery from circulation. The goal of an endograft is to prevent rupture or sac enlargement of an aneurysm by depressurizing the arterial walls.

Endografts are comparatively larger than stents and can require very large sheaths (>18F for endovascular aortic repair). To treat an aortic aneurysm with an endograft, there must be adequate sealing zones of healthy aorta above and below the aneurysm.



Please see the AAA Chapter for a full review of endova scular aortic repair (EVAR). Please see the TAA/TAAA Chapter for a full review of thoracic endova scualr aortic repair (TEVAR)

For patients that have extension of the aneurysm into the visceral vessels of the aorta, the use of a conventional, non-fenestrated endograft would block blood flow to the branches of the




Figure 5: Balloon-expandable stent deployment courtesty of Blausen Medical. Retrieved on 22 February 20. Wikimedia.

diseased segment of the aorta, such as the kidneys or bowel. Treatment of thoracoabdominal or complex abdominal aortic aneurysms with endovascular repair requires a graft that allows flow to branches of the aorta. Some common techniques for maintaining visceral blood flow include:

Fenestrated endografts include small, circular holes in the main body of the endograft. The french word for window is "fenêtre." When the fenstrations (or windows) are placed in line with visceral vessel origins, balloon-expandable stents can be used to bridge to the main endograft to the native renal or mesenteric arteries.

Additional characteristics about fenestrated grafts:

- They are ideal when the inner aortic diameter is small, making for a short distance (<5 mm) between the endograft and native visceral artery.
- They require the pathway between the endograft and the native artery to be close to perpendicular; that is, the pathway for the stent out of the graft cannot be at too large of an angle due to risk of stent fracture or kinking.
- The part of the stent inside the endograft must be flared out with a balloon to prevent endoleak from inside the endograft into the aneurysm sac.
- An of a commercially available fenestrated endograft is the Cook Medical Zenith Fenestrated (ZFen) AAA endograft. Click here to watch an animation of ZFen deployment.

Branched endografts have side branches built-into the main body of the graft in place of fenestration. These branches extend from the outer body of the main graft towards the target vessel.

Additional characteristics about branched grafts:

- They require a larger inner a ortic diameter for more space between the endograft and target artery.
- Branches can receive self-expanding or balloon-expandable stents.

Common Endovascular Procedures

Vascular surgeons perform a number of endovascular procedures. The most common include placement of an inferior vena cava (IVC) filter, diagnostic angiograms with angioplasty and stenting, fistulagrams, endovascular aneurysm repair (EVAR), and thrombectomy and thrombolysis. There are a number of basics that trainees should know before starting one of these procedures.

IVC Filter Placement

- An IVC Filter is a metal cage-like device placed in the IVC to prevent clots from embolizing to the pulmonary vasculature.
- The hook at the superior aspect of the filter is used to aid with retrieval.
- The struts will catch clots and will anchor into the walls of the IVC to prevent migration.
- Placement of an IVC filter can be controversial but it is commonly used to prevent pulmonary embolism in patients who have a strict contraindication to anticoagulation.
- They are sometimes placed prophylactically in high-risk trauma patients.




Figure 6: Inferior Vena Cava Filter courtesy of scientificanimations.com. Retrieved on 30 March. Wikimedia.

Procedure:

- Access is usually obtained via the right femoral vein as it gives direct access to the IVC.
- Ultrasound should be used to confirm that there is no thrombus at the access site Access is then obtained via the Seldinger technique and a venacavogram is then obtained to assess the anatomy and the location of the renal veins
- The filter is generally placed just below the renal veins to prevent renal vein thrombus if filter thrombosis occurs.
- If a retrievable filter is placed, it should be removed as soon as the filter is no longer clinically indicated.
- Filter retrieval becomes more challenging the longer the filter remains in place.
- Filter retrieval is generally performed through the internal jugular vein in the neck to give access to the hook at the superior end of the filter, which can be snared and removed.

Note

Please find a more detailed review of IVC filters here

Lower Extremity Angiogram

A lower extremity angiogram is the starting point for a vascular surgeon to perform a number of different interventions including angioplasty, stenting, and thrombectomy. An angiogram is a procedure in which contrast dye is used to display the vasculature and its disease burden under X-ray.



Tip

Before an angiogram, one should be aware of all prior interventions a patient has had, both open and endovascular. It can be useful to review old operative reports and imaging studies in order to know the patient's current anatomy.



🕊 Tip

In addition, the patient's most recent imaging studies should be reviewed including ABI/PVRs, arterial duplex, and any axial imaging such as a CTA.

Please see Claudication Chapter for a summary of ABI/PVR testing and interpretation.

Procedure:

- Access is typically obtained in the contralateral common femoral artery and a wire and catheter are passed up and over the aortic bifurcation to gain access to the operative (ipsilateral) side.
- A series of angiograms are taken to assess the level of disease and based on this a final operative plan will be made.
- There are multiple options for treatment including angioplasty, stenting, atherectomy, thrombectomy, and thrombolysis, to name a few.

Possible Interventions:

- Angioplasty: a balloon is advanced across the area of concern (usually atherosclerosis causing a flow limitation) and expanded to dilate the area of stenosis.
- Stenting: commonly after an angioplasty, a stent is placed to maintain patency of the area. There are multiple types of stents (covered above) that can be used depending on anatomic location, size, and indication.
- Atherectomy: a procedure where a calcified plaque is physically removed from the artery. This can be done with a cutting blade, a spinning drill-like device, or a laser.

Note

Considerations for Atherectomy

As plaque is broken into smaller pieces, there is a significant risk for **distal embolization** of plaque debris. To prevent this, an embolic protection device is often used which can filter blood and catch any smaller particulate matter that travels distally. Some of the devices include an aspiration device to remove clot and thrombus as it is removed. There is ongoing research into the outcomes of atherectomy versus stenting alone, but generally, the long-term patency rate is similar. ("Atherectomy for Lower Extremity Intervention: Why, When, and Which Device?" n.d.)

- Thrombolysis: a procedure where the clot is dissolved often with tPA and heparin to treat an acute thrombotic event. Often, angiograms are performed for acute limb ischemia from a clot rather than atherosclerotic disease. A lysis catheter or Cragg-Macnamara catheter can be placed. The Cragg-Macnamara catheter is a long catheter with multiple side ports where medication can be infused over the length of a thrombosed artery. This catheter can be left in place for 24-48 hours and the medication is allowed to "bust" the acute clot. After this, the patient will be brought back to the operating room to evaluate the progress of the clot dissolution. (Giannakakis et al. 2017)
- Thrombectomy: a procedure where acute or subacute thrombus is removed from a vessel. This can be accomplished in a variety of ways. Commonly, a suction thrombectomy system can be used to remove thrombus from a vessel using vacuum force for example using the **Penumbra system.** Alternatively, a mechanical thrombectomy can be performed where a Fogarty catheter is deployed past a thrombus, a balloon inflated at the end of the catheter, and then manually pulled back towards an arteriotomy and the clot then removed.(Karnabatidis et al. 2011)

Fistulograms

A fistulogram is an angiographic study used to interrogate and troubleshoot a disfunctional arteriovenous fistula.

Common complications of fistulas include:

- Anastomotic strictures.
- Inflow issues (such as atherosclerosis of the inflow artery).
- Outflow issues (such as central venous stenosis).
- Thrombosed fistula.

Access is obtained in different locations depending on the culprit lesions (the inflow artery, the fistula itself, the outflow veins) and can be treated with angioplasty and or stenting.

Risks

As with any invasive procedure, even a minimally invasive procedure comes with its own set of risks, and as the complexity of the procedure increases so does the risk.

The American Heart Association has put out guidelines to help stratify patients at risk for vascular complications.(Patel et al. 2010a)

Low risk cases were defined as those with a complication rate below 1%. Characteristics include:

- Diagnostic angiograms without interventions.
- Short operative times.
- Small sheaths (5 Fr).
- Minimal anticoagulation.
- Younger male patients with larger body habitus and normal renal function were at lowest risk of complications.

Moderate risk cases were defined as those with a complication rate of 1-3%. Risk factors include:

- Longer, more complex procedures.
- Involve larger sheaths (6-7Fr).

High risk cases were defined as those with a complication rate >3%. Risk factors include:

- Peripheral arterial disease patients
- Advanced age
- Female sex at birth
- Liver disease
- Coagulopathy
- Immunosuppression
- Status post valve replacement
- Renal dysfunction.
- Longer procedure times
- Larger sheaths (>8Fr), use of arterial closure devices
- More likely to involve anticoagulation

Access Complications

Although each procedure comes with its own specific risks (e.g. stroke for a carotid artery stent, distal embolization for atherectomy etc), there are multiple complications that can occur with any endovascular procedure. The procedure specific complications will be explored in further detail in each chapter. Common access complications are listed below.

Groin Hematoma

A groin hematoma is a collection of blood that has collected outside the lumen of the vessel, which often occurs after removal of the sheath.

Groin hematoma characteristics:

- Occur in about 1-3 % of cases
- Often due to a stick not directly over the femoral head in which case manual pressure may not provide adequate hemostasis.
- Hematomas are rarely ever operated on unless they case skin compromise, severe pain, femoral nerve compression, or are rapidly expanding.
- Hematoma will generally be absorbed by the body.
- If they are operated on, there is a significant risk of infection (about 20%) and seroma formation.

Pseudoaneurysms

A pseudoaneurysm is defined as a collection of turbulent blood flow contained only by one or two layers of the arterial wall as opposed to a true aneurysm where all three layers are dilated (occur in less than 1% of endovascular cases). They occur secondary to injury (iatrogenic or trauma).

These must be managed urgently and treated appropriately to prevent significant morbidity and mortality.

Pseudoaneurysm presentation:

- Pulsatile often painful mass at a prior access site
- May lead to skin necrosis
- Bruit on auscultation

Pseudoaneurysm risks:

- Vessel rupture
- Thrombus formation with distal embolization
- Skin necrosis
- Nerve compression
- Venous compression

Etiology:

Most often, seen at the femoral artery as an access site complication from:

- Failed deployment of a closure device
- Laceration of the artery or branch by an access needle
- Inadequate pressure or length of pressure being held.

Patients who are obese, female, hypertensive or are on anticoagulation or antiplatelet therapy are at higher risk for femoral PSA.

Workup:

- Duplex ultrasound of the site.
- Pulse exam to ensure adequate distal perfusion.

Management:

Management is dependent on size and presentation of the pseudoaneurysm. In the past, all pseudoaneurysms were treated operatively, but with new advances, minimally invasive techniques have become more common with good outcomes.

- For a small asymptomatic pseudoaneurysm (less than 2cm), an appropriate first step would be observation and repeat duplex ultrasound in about one month.
- For a persistent pseudoaneurysm after one month, a larger/loculated asymptomatic pseudoaneurysm, or a pseudoaneurysm that presents with minor pain at the site, minimally invasive intervention with ultrasound guided compression or ultrasound guided thrombin injection is appropriate.
- For any pseudoaneurysm presenting with persistent hemodynamic instability, limb ischemia, skin necrosis, AV fistula, nerve compression, expanding hematoma, an urgent operation is required.
- Urgent open repair generally consists of gaining proximal and distal control and then primary repair of the arterial defect.
- Urgent endovascular management may also be entailed by deploying coils in the sac and placing a covered stent in the femoral artery, therefore thrombosing the pseudoaneurysm.

[Kassem et al. (2013)](Kassem and Elmahdy 2014)(Tulla, Kowalski, and Qaja 2024)

Retroperitoneal Hematoma

A retroperitoneal hematoma is a collection of blood accumulating in the retroperitoneal space where the external iliac artery travels.

Key points:

- It is a rare complication occurring in about 0.5% of cases.
- Retroperitorally hematomas are a dreaded complication as they comes with high morbidity and mortality especially if not rapidly diagnosed.
- Retroperitoneal hematomas occur when the arterial stick in the artery during groin puncture is performed too high, above the level of the inferior epigastric artery.
- This leads to extravasation of blood into the retroperitoneal space after sheath removal.
- Retroperitoneal hematomas can usually can be treated with a closure device if diagnosed intra operatively or a covered stent from the contralateral side.

Arteriovenous (AV) Fistula

An arteriovenous fistula is an aberrant connection between an artery and a vein. Although commonly formed surgically for hemodialysis access, they can be caused introgenically during arterial access for endovascular intervention.

Key points:

- They are rare and occur in less than 1% of groin access cases.
- Occur during inadvertent puncture of the deep femoral artery and deep femoral vein.
- Usually no intervention is needed.

Dissection

An arterial dissection is the separation of the layers of the arterial wall.

Key points:

- May occur naturally (often secondary to hypertension) or introgenically (in the setting of arterial puncture). This most often occurs in the setting of a posteriorly lying plaque in the artery where the wire can separate the posterior layers.
- To prevent dissection, do not apply too much forward pressure when advancing a wire during initial arterial access. Do not inject fluid (saline or contrast) unless there is blood return to ensure the needle or catheter tip are in the true lumen.

• Treatment can be conservative in the case of an introgenic dissection or treated with stent placement if the dissection limits blood flow. Intravascular ultrasound can be used to further assess the dissection.

Acute Thrombosis of Access Vessel

Acute thrombus of the access vessel is when the artery where access is obtained completely "clots off." It is an incredibly rare complication caused by incorrect deployment of vascular closure devices, access site dissection, or occlusive sheath placement in a small or diseased vessel.

Endovascular Surgical Complications

Atheroembolism

Atheroembolism is the dislodgement of a portion of a cholesterol plaque leading to distal embolization causing acute thrombosis or partial occlusion.

Key points: (Liew and Bartholomew 2005)

- Generally, when these come from the ascending aorta or arch, they will embolize to the cerebral and retinal vessels. On the other hand, when an embolus originates in the thoracic or descending aorta, it will likely embolize to the mesenteric, renal or lower extremity vessels.
- These emboli will then cause a further inflammatory response worsening outcomes.
- Outcomes are dependent on volume and size of emboli and can range from sub-clinical to fatal
- Embolization is most common to the skin, kidneys, and GI tract though it is dependent on the site of the procedure.
- Unfortunately this condition is generally under diagnosed and challenging to treat. Primary prevention is of utmost importance with good surgical technique.

Wire Perforation

Wire perforation occurs when an angiographic wire creates a hole in a vessel.

Key points:

- Incredibly rare but treatable when diagnosed rapidly.
- Occurs when a large stiff hydrophilic wire is passed through a small vessel especially if not done under fluoroscopic guidance.
- Treatment involves transcatheter embolization of the perforation site.

Vessel Rupture

Vessel rupture occurs when a device or balloon is oversized for a vessel and causes a "blowout" of a vessel.

Key points:

- Rupture can be cause by an oversized balloon or balloon expandable stent, by expanding a properly sized balloon in a heavily calcified vessel or by passage of a large device such as an EVAR device through a small vessel.
- Rupture is associated with a high morbidity and mortality especially if there is any delay in diagnosis.
- Immediately upon recognition of vessel rupture, a balloon should be inflated in the region to obtain adequate hemostasis. Simultaneously, supplies should be can gathered for vessel repair.
- Repair is usually done with a covered stent placed across the rupture. If adequate hemostasis cannot be obtained, open repair may be necessary.

Teaching Case

Scenario

A 71 year old male with a past medical history of hypertension, hyperlipidemia, coronary artery disease with stent placement was diagnosed with acute type B aortic dissection three weeks ago. He was managed medically in the hospital and sent home after one week with blood pressure and pain well-controlled. An image from the CTA at time of diagnosis is seen below:

The proximal extent of the dissection was 3 cm distal to the left subclavian artery and the distal extent was 2 cm superior to the celiac artery, allowing good landing zones for an endograft. The left internal iliac artery is fully occluded, and the right internal iliac artery has roughly 50% stenosis. The left external iliac artery is heavily calcified and stenotic >50%. The right common and external iliac artery are patent without evidence of significant atherosclerosis. The patient is evaluated in the preoperative area prior to placement of a thoracic endograft in the hybrid operating room.



Figure 7: CTA

Exam

HEENT: Pupils round and reactive to light, no lymphadenopathy

Cardiac: Regular rate and rhythm

Pulmonary: Clear to auscultation in all lung fields. No accessory muscle use.

Abdominal: Soft, non-tender, non-distended, no pulsatile masses felt.

Vascular/Extremities: Warm, non-edematous. Palpable pulses in all four extremities.

Discussion Points

N.B. There is no prepared answers for the questions below.

However, we feel this chapter contains all the necessary information to answer the questions. If not, please let us know!

1. What clinical risk factors make this patient a good candidate for endovascular vs. open repair?

- 2. The endograft used for this operation requires at least an 18F sheath, which site of access should be used to insert and implant the graft?
- 3. What diameter and hydrophilicity of wire should be used to advance the thoracic endograft to the site of dissection?
- 4. The attending surgeon requested mean arterial pressures to be at least 90 mmHg throughout the operation. Why would intraoperative hypotension be concerning in this patient? What anatomic factors make this patient at particular risk?
- 5. If the patient developed pain at the site of access one week later, with a well-healing wound and palpable thrill on physical exam, what kind of access-related complication are you worried about and what would be your next step to confirm the diagnosis?

Key Articles

- Patel MR, Jneid H, Derdeyn CP, Klein LW, Levine GN, Lookstein RA, White CJ, Yeghiazarians Y, Rosenfield K; American Heart Association Diagnostic and Interventional Cardiac Catheterization Committee of the Council on Clinical Cardiology, Council on Cardiovascular Radiology and Intervention, Council on Peripheral Vascular Disease, Council on Cardiovascular Surgery and Anesthesia, and Stroke Council. Arteriotomy closure devices for cardiovascular procedures: a scientific statement from the American Heart Association. Circulation. 2010 Nov 2;122(18):1882-93. doi: 10.1161/CIR.0b013e3181f9b345. Epub 2010 Oct 4. Erratum in: Circulation. 2010 Nov 2;122(18):e507. PMID: 20921445. (Patel et al. 2010b)
- 2. Siracuse JJ, Farber A, Cheng TW, Raulli SJ, Jones DW, Kalish JA, Smeds MR, Rybin D, Schermerhorn ML; Vascular Quality Initiative. Common femoral artery antegrade and retrograde approaches have similar access site complications. J Vasc Surg. 2019 Apr;69(4):1160-1166.e2. doi: 10.1016/j.jvs.2018.06.226. Epub 2018 Dec 4. PMID: 30527937. (Siracuse et al. 2019)

Additional Resources

Audible Bleeding Content

- Holding Pressure Case Prep Endovascular Basics
- Audible Bleeding Exam Prep: Endovascular Access



• TeachMe Surgery: Pseudoaneurysm

Serious Games

Touch Surgery Simulations.

- Must download the Medtronic Touch Surgery mobile application to access the modules. Available for Apple and Android mobile devices.
- Femoral Artery Access

Gore Combat Manual

The Gore Medical Vascular and Endovascular Surgery Combat Manual is an informative and entertaining read intended as a vascular surgery crash course for medical students, residents, and fellows alike. Highly accessible with a thoughtfully determined level of detail, but lacking in learning activities (e.g. questions, videos, etc.), this resource is a wonderful complement to the APDVS eBook.



🕊 Tip

Please see pages 25-41, 49-61.

Operative Footage

Developed by the Debakey Institute for Cardiovascular Education & Training at Houston Methodist. YouTube account required as video content is age-restricted. Please create and/or log in to your YouTube account to have access to the videos.

Ultrasound-Guided Femoral Access

ProGlide Vascular Closure Device Deployment

Inferior Vena Cava (IVC) Filter Placement

Standard IVC Filter Retrieval

Anonymous Feedback

We welcome suggestions for updates. We want to hear from you! Please help us help you! To this end, please complete this anonymous Google Form to send us any feedback you may have.

Thank you for your continued support!



References

- AbuRahma, Ali F., Efthymios D. Avgerinos, Robert W. Chang, R. Clement Darling, Audra A. Duncan, Thomas L. Forbes, Mahmoud B. Malas, et al. 2022. "Society for Vascular Surgery Clinical Practice Guidelines for Management of Extracranial Cerebrovascular Disease." Journal of Vascular Surgery 75 (1): 4S-22S. https://doi.org/10.1016/j.jvs.2021.04.073.
- Allison, Matthew A., David G. Armstrong, Philip P. Goodney, Naomi M. Hamburg, Lee Kirksey, Kristie J. Lancaster, Carlos I. Mena-Hurtado, Sanjay Misra, Diane J. Treat-Jacobson, and Khendi T. White Solaru. 2023. "Health Disparities in Peripheral Artery Disease: A Scientific Statement From the American Heart Association." Circulation 148 (3): 286–96. https://doi.org/10.1161/cir.00000000000001153.
- Arya, Shipra, Sujin Lee, Greg J. Zahner, Beth E. Cohen, Jade Hiramoto, Owen M. Wolkowitz, Anjali Khakharia, Zachary O. Binney, and S. Marlene Grenon. 2018. "The Association of Comorbid Depression with Mortality and Amputation in Veterans with Peripheral Artery Disease." *Journal of Vascular Surgery* 68 (2): 536–545.e2. https://doi.org/10.1016/j.jvs.2017.10.092.
- Assari, Shervin. 2014. "Veterans and Risk of Heart Disease in the United States: A Cohort with 20 Years of Follow Up." *International Journal of Preventive Medicine* 5 (6): 703–9. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4085922/.
- "Atherectomy for Lower Extremity Intervention: Why, When, and Which Device?" n.d. https://www.acc.org/latest-in-cardiology/articles/2015/06/16/07/58/http%3a% 2f%2fwww.acc.org%2flatest-in-cardiology%2farticles%2f2015%2f06%2f16%2f07%2f58% 2fatherectomy-for-lower-extremity-intervention.
- Bazikian, Sebouh, Diego Urbina, Chiu-Hsieh Hsu, Karla A Gonzalez, Emily R Rosario, Dan I Chu, Jennifer Tsui, and Tze-Woei Tan. 2023. "Examining Health Care Access Disparities in Hispanic Populations with Peripheral Artery Disease and Diabetes." *Vascular Medicine* 28 (6): 547–53. https://doi.org/10.1177/1358863x231191546.
- Beckman, Joshua A., Peter A. Schneider, and Michael S. Conte. 2021. "Advances in Revascularization for Peripheral Artery Disease: Revascularization in PAD." *Circulation Research* 128 (12): 1885–1912. https://doi.org/10.1161/circresaha.121.318261.
- "Beneficial Effect of Carotid Endarterectomy in Symptomatic Patients with High-Grade Carotid Stenosis." 1991. New England Journal of Medicine 325 (7): 445–53. https://doi.org/10.1056/nejm199108153250701.
- Berchiolli, Raffaella, Giulia Bertagna, Daniele Adami, Francesco Canovaro, Lorenzo Torri, and Nicola Troisi. 2023. "Chronic Limb-Threatening Ischemia and the Need for Revascularization." *Journal of Clinical Medicine* 12 (7): 2682. https://doi.org/10.3390/jcm12072682.
- Blanchette, Virginie, Malindu E. Fernando, Laura Shin, Vincent L. Rowe, Kenneth R. Ziegler,

- and David G. Armstrong. 2022. "Evolution of WIfI: Expansion of WIfI Notation After Intervention." The International Journal of Lower Extremity Wounds, September, 153473462211228. https://doi.org/10.1177/15347346221122860.
- Chaikof, Elliot L., Ronald L. Dalman, Mark K. Eskandari, Benjamin M. Jackson, W. Anthony Lee, M. Ashraf Mansour, Tara M. Mastracci, et al. 2018. "The Society for Vascular Surgery Practice Guidelines on the Care of Patients with an Abdominal Aortic Aneurysm." *Journal of Vascular Surgery* 67 (1): 2–77.e2. https://doi.org/10.1016/j.jvs.2017.10.044.
- Conte, Michael S., Andrew W. Bradbury, Philippe Kolh, John V. White, Florian Dick, Robert Fitridge, Joseph L. Mills, et al. 2019a. "Global Vascular Guidelines on the Management of Chronic Limb-Threatening Ischemia." *Journal of Vascular Surgery* 69 (6): 3S–125S.e40. https://doi.org/10.1016/j.jvs.2019.02.016.
- ———, et al. 2019b. "Global Vascular Guidelines on the Management of Chronic Limb-Threatening Ischemia." *Journal of Vascular Surgery* 69 (6): 3S–125S.e40. https://doi.org/10.1016/j.jvs.2019.02.016.
- ———, et al. 2019c. "Global Vascular Guidelines on the Management of Chronic Limb-Threatening Ischemia." *Journal of Vascular Surgery* 69 (6): 3S–125S.e40. https://doi.org/10.1016/j.jvs.2019.02.016.
- ———, et al. 2019d. "Global Vascular Guidelines on the Management of Chronic Limb-Threatening Ischemia." *Journal of Vascular Surgery* 69 (6): 3S–125S.e40. https://doi.org/10.1016/j.jvs.2019.02.016.
- ———, et al. 2019e. "Global Vascular Guidelines on the Management of Chronic Limb-Threatening Ischemia." *Journal of Vascular Surgery* 69 (6): 3S–125S.e40. https://doi.org/10.1016/j.jvs.2019.02.016.
- Conte, Michael S., Frank B. Pomposelli, Daniel G. Clair, Patrick J. Geraghty, James F. McKinsey, Joseph L. Mills, Gregory L. Moneta, et al. 2015a. "Society for Vascular Surgery Practice Guidelines for Atherosclerotic Occlusive Disease of the Lower Extremities: Management of Asymptomatic Disease and Claudication." Journal of Vascular Surgery 61 (3): 2S-41S.e1. https://doi.org/10.1016/j.jvs.2014.12.009.
- ———, et al. 2015b. "Society for Vascular Surgery Practice Guidelines for Atherosclerotic Occlusive Disease of the Lower Extremities: Management of Asymptomatic Disease and Claudication." *Journal of Vascular Surgery* 61 (3): 2S–41S.e1. https://doi.org/10.1016/j.jvs.2014.12.009.
- Darling, R. Clement, and C. Keith Ozaki, eds. 2016. Vascular Surgery. Arterial Procedures. Master Techniques in Surgery. Philadelphia: Wolters Kluwer.
- Dillavou, Ellen D., Satish C. Muluk, and Michel S. Makaroun. 2006. "A Decade of Change in Abdominal Aortic Aneurysm Repair in the United States: Have We Improved Outcomes Equally Between Men and Women?" *Journal of Vascular Surgery* 43 (2): 230–38. https://doi.org/10.1016/j.jvs.2005.09.043.
- DiLosa, Kathryn L., Ryan Khoa Nguyen, Christina Brown, Aidan Waugh, and Misty D. Humphries. 2023. "Defining Vascular Deserts to Describe Access to Care and Identify Sites for Targeted Limb Preservation Outreach." *Annals of Vascular Surgery* 95 (September): 125–32. https://doi.org/10.1016/j.avsg.2023.05.025.
- Gerhard-Herman, Marie D., Heather L. Gornik, Coletta Barrett, Neal R. Barshes, Matthew

- ———, et al. 2017b. "2016 AHA/ACC Guideline on the Management of Patients With Lower Extremity Peripheral Artery Disease: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines." Circulation 135 (12). https://doi.org/10.1161/cir.0000000000000470.
- Giannakakis, Sotirios, George Galyfos, Ioannis Sachmpazidis, Kostas Kapasas, Stavros Kerasidis, Ioannis Stamatatos, Georgios Geropapas, Georgios Kastrisios, Gerasimos Papacharalampous, and Chrisostomos Maltezos. 2017. "Thrombolysis in Peripheral Artery Disease." Therapeutic Advances in Cardiovascular Disease 11 (4): 125–32. https://doi.org/10.1177/1753944716687517.
- Hackler, Eddie L., Naomi M. Hamburg, and Khendi T. White Solaru. 2021. "Racial and Ethnic Disparities in Peripheral Artery Disease." *Circulation Research* 128 (12): 1913–26. https://doi.org/10.1161/circresaha.121.318243.
- Hardman, Rulon, Omid Jazaeri, J. Yi, M. Smith, and Rajan Gupta. 2014. "Overview of Classification Systems in Peripheral Artery Disease." Seminars in Interventional Radiology 31 (04): 378–88. https://doi.org/10.1055/s-0034-1393976.
- "Health Disparities | DASH | CDC." 2023. https://www.cdc.gov/healthyyouth/disparities/index.htm.
- Hoel, Andrew W., Ahmed Kayssi, Soma Brahmanandam, Michael Belkin, Michael S. Conte, and Louis L. Nguyen. 2009. "Under-Representation of Women and Ethnic Minorities in Vascular Surgery Randomized Controlled Trials." *Journal of Vascular Surgery* 50 (2): 349–54. https://doi.org/10.1016/j.jvs.2009.01.012.
- Howard, Dominic P J, Liam Gaziano, and Peter M Rothwell. 2021. "Risk of Stroke in Relation to Degree of Asymptomatic Carotid Stenosis: A Population-Based Cohort Study, Systematic Review, and Meta-Analysis." *The Lancet Neurology* 20 (3): 193–202. https://doi.org/10.1016/s1474-4422(20)30484-1.
- Karnabatidis, Dimitris, Stavros Spiliopoulos, Dimitrios Tsetis, and Dimitris Siablis. 2011. "Quality Improvement Guidelines for Percutaneous Catheter-Directed Intra-Arterial Thrombolysis and Mechanical Thrombectomy for Acute Lower-Limb Ischemia." *Cardio-Vascular and Interventional Radiology* 34 (6): 1123–36. https://doi.org/10.1007/s00270-011-0258-z.
- Kassem, Hussein Heshmat, and Mahmoud Farouk Elmahdy. 2014. Femoral Artery Pseudoaneurysm Current Diagnosis and Treatment. LAP LAMBERT Academic Publishing. https://doi.org/10.13140/2.1.3471.2640.
- Kassem, Hussein Heshmat, Mahmoud Farouk Elmahdy, Essam Baligh Ewis, and Soilman Ghareeb Mahdy. 2013. "Incidence and Predictors of Post-Catheterization Femoral Artery Pseudoaneurysms." *The Egyptian Heart Journal* 65 (3): 213–21. https://doi.org/10.1016/j.ehj.2012.07.003.
- Kazis, Lewis E., Donald R. Miller, Jack Clark, Katherine Skinner, Austin Lee, William

- Rogers, Avron Spiro, et al. 1998. "Health-Related Quality of Life in Patients Served by the Department of Veterans Affairs." *Archives of Internal Medicine* 158 (6): 626. https://doi.org/10.1001/archinte.158.6.626.
- Khan, Tahir, Falahat Farooqui, and Khusrow Niazi. 2008. "Critical Review of the Ankle Brachial Index." Current Cardiology Reviews 4 (2): 101–6. https://doi.org/10.2174/157340308784245810.
- Lederle, Frank A., Tassos C. Kyriakides, Kevin T. Stroupe, Julie A. Freischlag, Frank T. Padberg, Jon S. Matsumura, Zhiping Huo, and Gary R. Johnson. 2019. "Open Versus Endovascular Repair of Abdominal Aortic Aneurysm." New England Journal of Medicine 380 (22): 2126–35. https://doi.org/10.1056/nejmoa1715955.
- Lederle, Frank A., Samuel E. Wilson, Gary R. Johnson, Donovan B. Reinke, Fred N. Littooy, Charles W. Acher, David J. Ballard, et al. 2002. "Immediate Repair Compared with Surveillance of Small Abdominal Aortic Aneurysms." *New England Journal of Medicine* 346 (19): 1437–44. https://doi.org/10.1056/nejmoa012573.
- Lee, Michael Ho-Yan, Pei Ye Li, Ben Li, Audrey Shakespeare, Yasith Samarasinghe, Tiam Feridooni, Cesar Cuen-Ojeda, Lina Alshabanah, Teruko Kishibe, and Mohammed Al-Omran. 2022. "A Systematic Review and Meta-Analysis of Sex- and Gender-Based Differences in Presentation Severity and Outcomes in Adults Undergoing Major Vascular Surgery." Journal of Vascular Surgery 76 (2): 581–594.e25. https://doi.org/10.1016/j.jvs.2022.02.030.
- Liew, Yin Ping, and John R Bartholomew. 2005. "Atheromatous Embolization." Vascular Medicine 10 (4): 309–26. https://doi.org/10.1191/1358863x05vm640ra.
- "Mapping Healthcare Deserts: 80." n.d. https://www.goodrx.com/healthcare-access/research/healthcare-deserts-80-percent-of-country-lacks-adequate-healthcare-access.
- McDermott, Mary M., and Michael H. Criqui. 2018. "Ankle-Brachial Index Screening and Improving Peripheral Artery Disease Detection and Outcomes." *JAMA* 320 (2): 143. https://doi.org/10.1001/jama.2018.8953.
- Mills, Joseph L., Michael S. Conte, David G. Armstrong, Frank B. Pomposelli, Andres Schanzer, Anton N. Sidawy, and George Andros. 2014. "The Society for Vascular Surgery Lower Extremity Threatened Limb Classification System: Risk Stratification Based on Wound, Ischemia, and Foot Infection (WIfI)." Journal of Vascular Surgery 59 (1): 220–234.e2. https://doi.org/10.1016/j.jvs.2013.08.003.
- Misra-Hebert, Anita D., Laura Santurri, Richard DeChant, Brook Watts, Michael Rothberg, Ashwini R. Sehgal, and David C. Aron. 2015. "Understanding the Health Needs and Barriers to Seeking Health Care of Veteran Students in the Community." Southern Medical Journal 108 (8): 488–93. https://doi.org/10.14423/SMJ.000000000000326.
- Parkinson, Fran, Stuart Ferguson, Peter Lewis, Ian M. Williams, and Christopher P. Twine. 2015. "Rupture Rates of Untreated Large Abdominal Aortic Aneurysms in Patients Unfit for Elective Repair." *Journal of Vascular Surgery* 61 (6): 1606–12. https://doi.org/10.1016/j.jvs.2014.10.023.
- Parodi, J. C., J. C. Palmaz, and H. D. Barone. 1991. "Transfemoral Intraluminal Graft Implantation for Abdominal Aortic Aneurysms." *Annals of Vascular Surgery* 5 (6): 491–99. https://doi.org/10.1007/bf02015271.
- Patel, Manesh R., Hani Jneid, Colin P. Derdeyn, Lloyd W. Klein, Glenn N. Levine, Robert

- A. Lookstein, Christopher J. White, Yerem Yeghiazarians, and Kenneth Rosenfield. 2010b. "Arteriotomy Closure Devices for Cardiovascular Procedures." *Circulation* 122 (18): 1882–93. https://doi.org/10.1161/cir.0b013e3181f9b345.
- ———. 2010a. "Arteriotomy Closure Devices for Cardiovascular Procedures." *Circulation* 122 (18): 1882–93. https://doi.org/10.1161/cir.0b013e3181f9b345.
- Qrareya, Mohammad, and Bara Zuhaili. 2021. "Management of Postoperative Complications Following Endovascular Aortic Aneurysm Repair." Surgical Clinics of North America 101 (5): 785–98. https://doi.org/10.1016/j.suc.2021.05.020.
- Remington, Patrick L, Bridget B Catlin, and Keith P Gennuso. 2015. "The County Health Rankings: Rationale and Methods." *Population Health Metrics* 13 (1). https://doi.org/10.1186/s12963-015-0044-2.
- Ricotta, John J., Ali AbuRahma, Enrico Ascher, Mark Eskandari, Peter Faries, and Brajesh K. Lal. 2011. "Updated Society for Vascular Surgery Guidelines for Management of Extracranial Carotid Disease: Executive Summary." *Journal of Vascular Surgery* 54 (3): 832–36. https://doi.org/10.1016/j.jvs.2011.07.004.
- Schaik, Theodorus G. van, Kak K. Yeung, Hence J. Verhagen, Jorg L. de Bruin, Marc R. H. M. van Sambeek, Ron Balm, Clark J. Zeebregts, et al. 2017. "Long-Term Survival and Secondary Procedures After Open or Endovascular Repair of Abdominal Aortic Aneurysms." Journal of Vascular Surgery 66 (5): 1379–89. https://doi.org/10.1016/j.jvs.2017.05.122.
- Schanzer, Andres, and Gustavo S. Oderich. 2021. "Management of Abdominal Aortic Aneurysms." New England Journal of Medicine 385 (18): 1690–98. https://doi.org/10.1056/NEJMcp2108504.
- Siracuse, Jeffrey J., Alik Farber, Thomas W. Cheng, Stephen J. Raulli, Douglas W. Jones, Jeffrey A. Kalish, Matthew R. Smeds, Denis Rybin, and Marc L. Schermerhorn. 2019. "Common Femoral Artery Antegrade and Retrograde Approaches Have Similar Access Site Complications." *Journal of Vascular Surgery* 69 (4): 1160–1166.e2. https://doi.org/10.1016/j.jvs.2018.06.226.
- "Social Determinants of Health: Key Concepts." n.d. https://www.who.int/news-room/questions-and-answers/item/social-determinants-of-health-key-concepts.
- Tulla, Katrina, Andrew Kowalski, and Erion Qaja. 2024. "Femoral Artery Pseudoaneurysm." In. Treasure Island (FL): StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK493210/.
- Walker, Michael D. 1995. "Endarterectomy for Asymptomatic Carotid Artery Stenosis." *JAMA: The Journal of the American Medical Association* 273 (18): 1421. https://doi.org/10.1001/jama.1995.03520420037035.
- Wijnand, Joep G. J., Devin Zarkowsky, Bian Wu, Steven T. W. van Haelst, Evert-Jan P. A. Vonken, Thomas A. Sorrentino, Zachary Pallister, et al. 2021. "The Global Limb Anatomic Staging System (GLASS) for CLTI: Improving Inter-Observer Agreement." *Journal of Clinical Medicine* 10 (16): 3454. https://doi.org/10.3390/jcm10163454.
- Willey, James, Amgad Mentias, Mary Vaughan-Sarrazin, Kimberly McCoy, Gary Rosenthal, and Saket Girotra. 2018. "Epidemiology of Lower Extremity Peripheral Artery Disease in Veterans." *Journal of Vascular Surgery* 68 (2): 527–535.e5. https://doi.org/10.1016/j.jvs.2017.11.083.